

**DIRECTIONS FOR MIXING**

Fill the spray tank about one-third full of water, start the agitator and pour in the required amounts of Supreme Oil. If other spray materials are to be used with the oil, first mix the other materials thoroughly with the water in the spray tank, and then add the oil last as the filling of the tank is being completed.

**DIRECTIONS FOR USE**

GALLONS INDICATED ARE AMOUNTS TO USE PER 100 GALLONS OF WATER.

**APPLES, PEARS - European**

Pract. Dormant Scale, San Jose Scale - Apply 2 gallons in the delayed dormant stage. Fruit Tree Leaf Hopper, Sawfly Scale - Apply 3 gallons in the delayed dormant stage. European Red Mite - Apply 2 gallons in the green tip to delayed dormant stage. Pear Leaf Blister Mite - Apply 3 gallons in the dormant stage. Pear Psylla - Apply 2 gallons in the late dormant stage.

Grasshopper, European Red Mite, San Jose Scale, Pear Leaf Blister Mite, Pear Psylla - Apply in the dormant stage. Pear Leaf Blister Mite - Apply 2 gallons in the delayed dormant stage.

**APPLES, PEARS - American**  
 Pear Leaf Blister Mite - Apply 2 gallons in the dormant stage. Pear Psylla - Apply 2 gallons in the dormant stage. Pear Leaf Blister Mite - Apply 2 gallons in the dormant stage. Pear Psylla - Apply 2 gallons in the dormant stage.

**PLUMS, PEACHES**  
 Pear Leaf Blister Mite - Apply 2 gallons in the dormant stage. Pear Psylla - Apply 2 gallons in the dormant stage. Pear Leaf Blister Mite - Apply 2 gallons in the dormant stage. Pear Psylla - Apply 2 gallons in the dormant stage.

**SUPREME OIL**

INSECTICIDE



ACCEPT

**ACTIVE INGREDIENT:**

Petroleum Oil (Superior Type) 97.0%

**INERT INGREDIENTS:** 3.0%

100.0%

**Petroleum Oil Specifications:**

Maximum Acid Residue (M.A.R.) 0.02%

Viscosity 134-136 ccs

**CAUTION**

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed

DO NOT STORE AT TEMPERATURES BELOW 32° F.

DO NOT USE IN UNDILUTED FORM.

AVOID INHALATION OF VAPORS AND SPRAY MIST

Caution: This is a flammable liquid. It is to be used in the spray tank top, and it is to be kept out of the reach of children and accepted for use in the spray tank.

MADE IN U.S.A.

REGISTERED TRADE MARK OF THE MANUFACTURER

© 1954 BY THE MANUFACTURER

100 GALLONS

Gallons

**DIRECTIONS FOR USE**

When a gun has been shut off or the outfit stopped for any length of time, spray on the ground for a few moments until the spray appears creamy white. This is desirable also where there are dead ends in the pipe lines of stationary outfits. All the material in stationary outfits should be sprayed out each day.

Spray on warm days in late Winter or early Spring, and before the buds separate into the clusters. Do not spray in freezing weather, or apply strong dosages after the buds have burst to any extent. Do not apply when buds have separated in the clusters.

Do not apply within 2 weeks of a sulfur spray or within 7 days of Captan.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milky white spray mixture results, provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank, with the agitator running, does not turn white, do not use it for either spray or the remainder of the product.

Do not use if the label on the product is only

DO NOT USE ON PLANTS OR MATERIALS WHICH ARE KNOWN TO BE SENSITIVE TO OILS OR TO WHICH A MIST OF OIL MAY BE HARMFUL. IN ACCORDANCE WITH FEDERAL REGULATIONS, THIS PRODUCT IS NOT TO BE USED ON PLANTS OR MATERIALS WHICH ARE KNOWN TO BE SENSITIVE TO OILS OR TO WHICH A MIST OF OIL MAY BE HARMFUL.

100