

MIXING DIRECTIONS: Mix 1 gallon of Contact Weedkiller in 50 to 60 gallons of water and spray to thoroughly wet all weed growth. Good tank agitation is desired during spraying operation, but not necessary.

GENERAL WEED CONTROL: Contact Weedkiller will control many common herbaceous weeds and grasses. It works best in warm weather. Perennials must be re-treated as new growth develops.

WEEDS & GRASSES CONTROLLED: Annual Bluegrass, Watergrass, Goosegrass, Redroot Pigweed, Lambsquarter, Kochia and Ragweed.

PRE-EMERGENCE WEED CONTROL: on cropland planted to BEANS, CORN, and POTATOES. Spray small weeds at least one day before emergence of the crop. Use up to 22½ gals. of diluted spray per acre. Do not apply to light soil or to shallow planted corn.

DRAINAGE DITCHBANKS & FENCEROWS: For the control of annual weeds and grasses use 1 gal. to 100 gals. of water and wet to point of run-off.


ORCHARD FLOORS: APPLES, APRICOTS, CHERRIES, PEARS, PEACHES, PLUMS, and PRUNES. BANKS OF DRAINAGE CANALS. Use up to 43 gallons of diluted spray per acre on crop areas. Apply a course spray thoroughly wetting all weed foliage. Apply directed spray to weed ground cover, and keep spray off foliage of the fruit trees. Destroy any fruit accidentally sprayed. Do not allow livestock to graze on treated ground cover.

PREHARVEST DESICATION OF ALFALFA, CLOVER and PEAS GROWN FOR SEED: Use 2 to 3 quarts of Contact Weedkiller per acre. Mix with 10 gallons of water for airplane applications and 30 gallons of water for ground sprayers. Apply 3-6 days before expected harvest. Do not graze treated areas or feed treated forage to livestock.

KILLING POTATO VINES: Use ½ gallon per acre 10-14 days before harvest. Degree and length of time for kill are determined by variety and vigor of vines and weather conditions at time of application. AIRPLANE APPLICATION: Use ½ gal

3468-10

ACCEPTED  
DECEMBER 2 1972  
UNDER THE GENERAL  
FUNGICIDE RODENT  
FOR ECONOMIC CONTROL  
ED UNDER THE  
TO ATTACK



# CONTACT WEEDKILLER

ACTIVE INGREDIENT:  
4,6-Dinitro-o-sec-butylphenol . . . . . 27%  
INERT INGREDIENTS: . . . . . 73%

## DANGER POISON



KEEP OUT OF REACH OF CHILDREN

**NOTE:** Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin or clothing. Do not take internally, avoid breathing spray drift. Do not wear contaminated clothing or shoes.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

EPA Reg. No. 3468-10

Manufactured by  
**SCHALL CHEMICAL, INC.** Monte Vista, Colorado

Mix 1 gallon of Contact Weedkiller in 50 to 60 gallons of water and spray to thoroughly wet all weed growth. Good tank agitation is desired during spraying operation, but not necessary.

Contact Weedkiller will control many common herbaceous weeds and grasses. It works best in warm weather. Perennials must be re-treated as new growth develops.

Annual Bluegrass, Watergrass, Goosegrass, Redroot Pigweed, Lambsquarter, Kochia and Ragweed.

on cropland planted to BEANS, CORN, and POTATOES. Spray small weeds at least one day before emergence of the crop. Use up to 22½ gals. of diluted spray per acre. Do not apply to light soil or to shallow planted corn.

For the control of annual weeds and grasses use 1 gal. to 100 gals. of water and wet to point of run-off.

APPLES, APRICOTS, CHERRIES, PEARS, PEACHES, PLUMS, and PRUNES. BANKS OF DRAINAGE CANALS. Use up to 43 gallons of diluted spray per acre on crop areas. Apply a course spray thoroughly wetting all weed foliage. Apply directed spray to weed ground cover, and keep spray off foliage of the fruit trees. Destroy any fruit accidentally sprayed. Do not allow livestock to graze on treated ground cover.

Use 2 to 3 quarts of Contact Weedkiller per acre. Mix with 10 gallons of water for airplane applications and 30 gallons of water for ground sprayers. Apply 3-6 days before expected harvest. Do not graze treated areas or feed treated forage to livestock.

Use ½ gallon per acre 10-14 days before harvest. Degree and length of time for kill are determined by variety and vigor of vines and weather conditions at time of application. **AIRPLANE APPLICATION:** Use ½ gal-

3468-10

**ACCEPTED**  
DECEMBER 2 1972  
UNDER THE FEDERAL FUNGICIDE ACT FOR ECONOMIC CONTROL TO ATTACH

# CONTACT WEEDKILLER

ACTIVE INGREDIENT:  
4,6-Dinitro-o-sec-butylphenol . . . . . 27%  
INERT INGREDIENTS: . . . . . 73%

DANGER  
POISON



**KEEP OUT OF REACH OF CHILDREN**

**NOTE:** Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin or clothing. Do not take internally, avoid breathing spray drift. Do not wear contaminated clothing or shoes.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

EPA Reg. No. 3468-10

Schall

# CONTACT WEEDKILLER

ACTIVE INGREDIENT:  
4,6-Dinitro-o-sec-butylphenol .....27%  
INERT INGREDIENTS: ..... 73%

## DANGER POISON



**KEEP OUT OF REACH OF CHILDREN**

**NOTE:** Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin or clothing. Do not take internally, avoid breathing spray drift. Do not wear contaminated clothing or shoes.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

EPA Reg. No. 3468-10

Manufactured by

**SCHALL CHEMICAL, INC.** Monte Vista, Colorado

lon with 9 1/2 gallons of water per acre. **GROUND APPLICATION:** Use 1/2 gallon in 50 gallons of water and spray to thoroughly wet all weed growth. Complete coverage of all parts of vines is essential. Do not graze livestock in treated areas.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, ponds, and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment, or disposal of wastes. Apply this product only as specified on this label.

### FIRST AID TREATMENT-ANTIDOTE

**SYMPTOMS OF POISONING.** Excessive Fatigue, Sweating, Thirst and Fever. If symptoms of poisoning develop from any type of exposure, **SEND FOR A PHYSICIAN.**

**FIRST AID:** Have patient lie quiet in coolest spot available. If feverish, cool with cold compresses or by immersion in cool water.

**IF SWALLOWED, SEND FOR A PHYSICIAN.** Induce vomiting by giving an emetic such as 2 tablespoons of table salt in a glass of warm water; repeat until vomit fluid is clear, then give two tablespoons of baking soda in a glass of water. Treat as in **FIRST AID**, above. If splashed in eyes, immediately flush eyes with plenty of water for at least 15 min. and get medical attention. If spilled on skin immediately remove contaminated clothing, including shoes, and wash skin with soap and plenty of water. If symptoms of poisoning develop, send for a physician and treat as in **FIRST AID**, above. Discard contaminated clothing and shoes, or clean them thoroughly before re-use. **NOTE TO ATTENDING PHYSICIAN:** Active ingredient is a metabolic stimulant. Treat symptomatically.

**NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

3468-10  
**ACCEPTED**  
DE 2 1972  
UNDER THE GENERAL FUNGICIDE FOR ECONOMIC TO ATTACK  
POISON  
DANGER  
POISON