

OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not contaminate water by cleaning equipment or disposal of waste and containers.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues will not exceed established tolerances, use only at recommended rates.

DIRECTIONS FOR USE

SPRAYING

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to animals may result. Do not combine with other materials.

LIVESTOCK OILERS OR BACKRUBBERS

For control of horn flies and as an aid to reduce lice infestations on beef cattle and swine, dilute this material at the rate of 1 gallon to 10 of diesel fuel. Place the mixture in oil reservoir or on burlap or oiler device and keep wet at all times. Place livestock oiler equipment where animals have free access to rubbing. Primarily near watering facilities or wherever animals congregate. Do not permit beef cattle and swine access to treatment within 30 days of slaughter or beef cattle and swine under 6 months of age. Do not use on dairy animals.

RESIDUAL SPRAY

For control of House Flies, Mosquitoes and Stable Flies. Use 1 gal. of this material to 9 gals. of water. Apply spray at the rate of 1/2 to 1 gal. per 1000 sq. ft. of surface. Thoroughly treat exteriors of barns, stock sheds and other farm buildings as well as manure, vegetation and refuse. Do not use on dairy barns or poultry houses. Repeat as needed.



1 GAL. NET

LIVESTOCK SPRAY, DIP & OILER CONC.

For Use on Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange and Ticks.

Contains .18 Lbs. Lindane and 4 Lbs. Toxaphene Per Gallon

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride (from Lindane)..... 1.9%
Toxaphene (Technical chlorinated camphene)*..... 42.0%
Xylene range aromatic solvent..... 49.7%

INERT INGREDIENTS:

*Chlorine Content 67% to 69%
100%

WARNING

Keep Out of Reach of Children

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not contaminate feed or foodstuffs. In case of contact, wash immediately with soap and water. Do not use on household pets or humans.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME
Protect From Temperatures Below 0°F.

USDA Reg. 3468-5

Manufactured by

SCHALL CHEMICAL, INC. Monte Vista, Colorado

ACCEPTED
2-2-68
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
OR ECONOMIC POISON REGISTER-
ED UNDER NO. *Comments 3468-12*

RECOMMENDATIONS

Pest	50 Gal. of Water	5 Gal. of Water
Beef Cattle	Amount of Concentrate to use Per:	
Horn Flies, Lice,	2 qt. for Sprays	1/2 cup
Sarcoptic Mange, Ticks	2 qt. for Dips	1/2 cup
Swine		
Lice, Sarcoptic Mange	2 qt. for Sprays	1/2 cup
	2 qt. for Dips	1/2 cup
Sheep		
Lice, Keds (Sheep Tick),		
Psoroptic or Sarcoptic Sheep	2 qt. for Sprays	1/2 cup
Scabs, Ticks	1 qt. for Dips	1/2 cup

APPLICATION

Horn Flies—spray back of animals every 3 weeks.
Lice, Ticks, Sarcoptic Mange and Sarcoptic Sheep Scab—immerse or spray animals thoroughly. Repeat after 2-3 weeks if needed.

CAUTION

Do not treat sick animals. Do not use on dairy animals or animals under 6 months of age. Do not spray animals within 30 days of slaughter or dip within 60 days of slaughter. Do not treat emaciated or lactating animals. Animals should have access to the shade following dipping or spraying in warm weather. Do not spray when weather is excessively cold or hot. Do not re-use empty container. Destroy container when empty.

NOTICE: Seller guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risks to persons or property rising out of the use or handling and accepts the product on these conditions.

ACCEPTED WITH COMMENTS

8

OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not contaminate water by cleaning equipment or disposal of waste and containers.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues will not exceed established tolerances, use only at recommended rates.

DIRECTIONS FOR USE

SPRAYING

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to animals may result. Do not combine with other materials.

LIVESTOCK OILERS OR BACKRUBBERS

For control of horn flies and as an aid to reduce lice infestations on beef cattle and swine, dilute this material at the rate of 1 gallon to 10 of diesel fuel. Place the mixture in oil reservoir or on burlap or oiler device and keep wet at all times. Place livestock oiler equipment where animals have free access to rubbing. Primarily near watering facilities or wherever animals congregate. Do not permit beef cattle and swine access to treatment within 30 days of slaughter or beef cattle and swine under 6 months of age. Do not use on dairy animals.

RESIDUAL SPRAY

For control of House Flies, Mosquitoes and Stable Flies. Use 1 gal. of this material to 9 gals. of water. Apply spray at the rate of 1/2 to 1 gal. per 1000 sq. ft. of surface. Thoroughly treat exteriors of barns, stock sheds and other farm buildings as well as manure, vegetation and refuse. Do not use on dairy barns or poultry houses. Repeat as needed.



1 GAL. NET

LIVESTOCK SPRAY, DIP & OILER CONC.

For Use on Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange and Ticks.

Contains .18 Lbs. Lindane and 4 Lbs. Toxaphene Per Gallon

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride (from Lindane).....	1.9%
Toxaphene (Technical chlorinated camphene)*.....	42.0%
Xylene range aromatic solvent.....	49.7%

INERT INGREDIENTS:

*Chlorine Content 67% to 69%	6.4%
	100%

WARNING

Keep Out of Reach of Children

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not contaminate feed or foodstuffs. In case of contact, wash immediately with soap and water. Do not use on household pets or humans.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME
Protect From Temperatures Below 0°F.

USDA Reg. 3468-5

Manufactured by
SCHALL CHEMICAL, INC. Monte Vista, Colorado

ACCEPTED
2-2-68

UNDER THE FEDERAL INSECTICIDE
HERBICIDE AND RODENTICIDE ACT
OR ECONOMIC POISON REGISTERED
UNDER NO. *Comments 3468-12*

RECOMMENDATIONS

Pest	50 Gal. of Water	5 Gal. of Water
Beef Cattle	Amount of Concentrate to use Per:	
Horn Flies, Lice,	2 qt. for Sprays	1/2 cup
Sarcoptic Mange, Ticks	2 qt. for Dips	1/2 cup
Swine		
Lice, Sarcoptic Mange	2 qt. for Sprays	1/2 cup
	2 qt. for Dips	1/2 cup
Sheep		
Lice, Keds (Sheep Tick),		
Psoroptic or Sarcoptic Sheep	2 qt. for Sprays	1/2 cup
Scabs, Ticks	1 qt. for Dips	1/2 cup

APPLICATION

Horn Flies—spray back of animals every 3 weeks.
Lice, Ticks, Sarcoptic Mange and Sarcoptic Sheep Scab—immerse or spray animals thoroughly. Repeat after 2-3 weeks if needed.

CAUTION

Do not treat sick animals. Do not use on dairy animals or animals under 6 months of age. Do not spray animals within 30 days of slaughter or dip within 60 days of slaughter. Do not treat emaciated or lactating animals. Animals should have access to the shade following dipping or spraying in warm weather. Do not spray when weather is excessively cold or hot. Do not re-use empty container. Destroy container when empty.

NOTICE: Seller guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risks to persons or property arising out of the use or handling and accepts the product on these conditions.

ACCEPTED WITH COMMENTS