

DIRECTIONS FOR USE

TIME OF APPLICATION - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.

SMALL GRAIN CROPS - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.


CORN - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.

SORGHUM - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.

PASTURES - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.

WOODY PLANT CONTROL - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.

GENERAL WEED CONTROL - Apply this product to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth. Apply to weeds in the early stages of their growth.



LV-4

WEED KILLER

(Low Volatile Emulsifiable)

ACTIVE INGREDIENTS:

2-Ethylhexyl (Isocetyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	69.2%
INERT INGREDIENTS:	30.8%
TOTAL	100.0%

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 45.9

Contains 4 lbs. 2,4 D Acid Equivalent Per Gallon

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN.
HARMFUL IF SWALLOWED. See Side Panels For Other Cautions.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

NET CONTENTS GALS

Manufactured for

SCHALL CHEMICAL, INC. Monte Vista, Delta, Colorado

ACCEPTED

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION UNDER NO. 3463-17

GENERAL INFORMATION

PREPARATION OF THE SPRAY - Fill the spray tank with half the required quantity of water, then add the recommended amount of LV-4 with agitation and continue filling the spray tank with water. The amount of water required in low volume applications may vary from 5 to 25 gallons per acre. For high volume applications 50 to 100 gallons of water or more may be necessary for good coverage. In any case use the same amount of 2,4 D recommended per acre.

CAUTION

Avoid contact with skin, eyes or clothing. Harmful if swallowed. In case of eye contact, wash with plenty of water. Avoid spray drift to susceptible plants.

DO NOT allow spray to drift to susceptible plants such as fruit trees, ornamentals, flowers, grapes, fruit trees and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays may cause leaf drop. Although this ester is much less volatile than butyl ester, spray drift of LV-4 at high temperatures may produce drift. The application which might cause LV-4 to spray plants is wind drift.

DO NOT use more than 1 gallon of this product per acre for most purposes.

DO NOT use the same spray equipment for applying other materials to plants as that with which this product is applied.

DO NOT use on sensitive plants, ornamentals, or trees.

DO NOT use on plants which are sensitive to phenoxyacetic acids.

DO NOT use on plants which are sensitive to phenoxyacetic acids.

WARRANTY AND LIMITATION OF DAMAGES

This product is warranted to be free from adulteration and to conform to the label directions. It is sold as a weed killer and is not intended for use as a herbicide. The user assumes all responsibility for the proper use of this product. The manufacturer is not liable for any damage to crops or property caused by the use of this product.