



# AMINE-4 Weed Killer

**ACTIVE INGREDIENTS:**  
 Dimethylamine Salt of 2,4-Dichloro-  
 phenoxyacetic Acid \* ..... 46.3%

**INERT INGREDIENTS:** ..... 53.7%

**TOTAL** ..... 100%

Each gallon contains 5.3 lbs. 2,4-D acid  
 Soluble in Water

\* Equivalent to 2,4-D chlorophenoxyacetic Acid 38.4%

**ISOMER SPECIFIC BY ADAC METHOD NO. 6 D01-5**

**CAUTION: KEEP OUT OF THE REACH OF CHILDREN. See Side Panels for Other Cautions.**

NET CONTENTS \_\_\_\_\_ GALS.

Manufactured by  
**SCHALL CHEMICAL, INC. Monte Vista, Colorado**  
CBI OFFSET PRINTERS Monte Vista CO

**SUSCEPTIBLE WEEDS**  
Some weeds listed below as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of crop.



**GENERAL INFORMATION**  
**PREPARATION OF THE SPRAY:** Fill the spray tank with half the required amount of water, then add the recommended amount of AMINE-4 with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage.

**CAUTIONS**  
Avoid contact with eyes, skin and clothing. Harmful if swallowed. May cause skin irritation. Do not contaminate irrigation ditches or water used for domestic purposes. Do not reuse empty container. Destroy by burying in a safe place. Do not burn.  
Although AMINE-4 is non volatile, care must be used to prevent spray drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other desirable plants sensitive to 2,4-D. Coarse sprays are especially likely to drift.  
**DO NOT** use the same spray equipment for applying other materials to plants, as injury may result.  
**DO NOT** store near fertilizers, feeds, seed, fungicides, or insecticides.  
**DO NOT** apply to lawns containing Bent, St. Augustine and Carpet grasses or Dichondra and Clover as injury may result.

**WARRANTY AND LIMITATION OF DAMAGES:**  
SELECH warrants that the material is for the uses, purposes and conditions for which it is intended and that the material is free from defects in material and workmanship under normal conditions of use. SELECH makes no other express or implied warranty, including any warranty of fitness for a particular purpose, and no agent of SELECH is authorized to do so. No written or printed reference to this warranty, in no event shall SELECH'S liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.  
EPA - Est 3468-CO-1 EPA REG. NO. 3468-20

## DIRECTIONS FOR USE

**TIME OF APPLICATION:** Best results are obtained when AMINE-4 is used on weeds that are young and in a rapid growing condition.

**SMALL GRAIN CROPS** (wheat, oats, rye): Apply AMINE-4 at the rate of 1/2 to 1 pint per acre on small grains when fully tilled or stubbed (4 to 8 inches tall) but before head emerges from the boot. Injury will be likely to occur if this product is applied earlier than "tiller" or later than "boot". Do not use on grains undersown with legumes such as alfalfa or clovers. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D.

**APPLICATION:** Aerial application should be mixed with water at the rate of 3 to 10 gallons per acre. Ground Application should be mixed with water at the rate of 10 to 25 gallons per acre. The specific amount of chemical to be used can be found under crop recommendations listed below.

**CORN:** For post-emergence, apply AMINE-4 at the rate of 1/2 to 1 pint per acre. When spraying corn over 10 inches in height, use drop nozzles directing the spray down to the base of the corn plant. Do not apply from tasseling to dough stage.

**SORGHUM:** Use spray at the rate of 1/2 to 1 pint of AMINE-4 per acre on plants 6 to 12 inches high when secondary roots are well established. Use drop nozzles when Sorghum is over 10 inches high directing the spray down to the base of the plant.

**PASTURE:** To control many weeds in pastures, meadows, use 1/2 pints per acre of AMINE-4. DO NOT apply to recently seeded pastures until grass is well established. Most legumes are usually injured or killed at the rates recommended. Do not graze treated area to dairy animals within 7 days after treatment.

**LAWNS:** Apply 2 to 3 pints of AMINE-4 per acre in sufficient water to obtain uniform coverage. Do not apply to Bent grass, Dichondra, St. Augustine or to freshly seeded turf. If grasses have become well established, Golf courses use above rates also.

**ROADSIDE AND VACANT LOT SPRAYING:** Use from 1 to 2 pints of AMINE-4 per acre, depending on the size of the weeds. Use sufficient water to give uniform coverage.

**NOTE WHEN STORED AT TEMPERATURES BELOW FREEZING:** It may be necessary to warm up to at least 45 degrees and shake thoroughly before using.