Sevin* - Sulfur

ACTIV	E INGRED																																	
	*Carbaryl	(1-n)	aphthy	1 N-	methy	ylca	rba	ma	te).	•		•		•		•	• •	•	••	•		•	•		•		•		•			•	5.	07
	Sulfur				• • •					•		•		•				•		•		•	•		•		•		•	•		•	.27.8	17
INERT	INGREDII	ENTS	: <i>.</i>							•						•		•		•							•		•	•		•	.67.1	7
*Sevin,	U.S. Pat.	No.	2,903,	478,	3,009	,855	•	T	rad	e- 11	nark	t of	ľ	nioı	a (Carl	obic	e C	orp	OT	lti (on	for	th	e a	ıcti	ve	in	zre	di€	ent	ca	ırbary	/1.

Caution: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

DISPOSAL WARNING: Do not reuse this container. Completely empty the contents. Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DIRECTIONS FOR USE: Repeat application as necessary. IMPORTANT read the entire label and conditions of sale including the Warranty before using 5 Sevin 50 Sulfur. Apply dust when insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. 5 Sevin 50 Sulfur may be applied up to the day of harvest. POTATOES: Colorado Potato Beetle, Flea Beetle, Leafhopper, Fall Armyworms, Stink Bugs, Tarnished Plant Bug — Use 20 to 40 lbs. per acre.

WARRANTY NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured by SCHALL CHEMICAL INC.

Monte Vista and Delta, Colorado EPA Reg. No. 3468-21

ACCEPTED WITH COMMENTS