

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing of dust. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Long pants, long sleeve shirt, gloves, hat and boots must be worn during mixing and loading.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in cool, dry location. Do not store in direct sunlight. Keep away from fire or sparks. Never allow Maneb to become wet during storage. This may lead to certain chemical changes which will reduce effectiveness of Maneb as a fungicide and create vapors which may be harmful. Keep container closed when not in use. Use dust mixture as soon as possible after preparation.

PESTICIDE DISPOSAL: Pesticide spray mixture or waste water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary manner or by incineration if allowed by State and local authorities.

MANEB-SULFUR

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebis(dithiocarbamate)) 6.0%
Manganese equivalent as metallic (23%) 1.23%
Sulfur 50.0%

INERT INGREDIENTS:

44.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed, induce vomiting with a mixture of soap or mustard in warm water. Call a physician immediately. If on skin, wash immediately with soap and water.

See Side Panel for additional Precautionary Statements.

EPA Reg. No. 3468-23 EPA Est. No. 3468-CO 1

Manufactured by SCHALL CHEMICAL, INC.
Monte Vista, Colorado 81144

DIRECTIONS FOR USE

It is a violation of the Federal law to use this product in a manner inconsistent with its labeling. Do not use this product at higher than recommended rate.


Apply dust when diseases first appear and repeat at intervals as required to upper and lower leaf surfaces. Observe day intervals between last application and harvest. Cantaloupe; anthracnose, cercospora leaf spot, downy mildew, gummy stem blight. Use 30 lbs. per acre. Begin at late bunching when vines begin to run or when disease first appears. Repeat at 7-10 day intervals. Do not apply within 5 days of harvest. Lettuce (Head), downy mildew. Use 30-40 lbs. per acre. Begin when disease appears. Repeat at 7-10 day intervals. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming. Potatoes, early and late blight. Use 30 lbs. per acre. Begin when plants are 2 to 6 inches high. Repeat at 5 to 10 day intervals, more often under severe late blight conditions. No time limitations.

NOTE: This pesticide is to be sold ONLY in its original unopened package.

**NOTICE — READ CAREFULLY
LIMITED WARRANTY STATEMENT**

Schall Chemical, Inc. warrants that this production conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Schall Chemical, Inc. neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material, in the event of any breach by Schall Chemical, Inc. of this limited warranty. Schall Chemical, Inc. shall not be liable for consequential damages.

Colson Printing Co. 14556



BEST DOCUMENT AVAILABLE