

# CYTHION\* SULFUR

**ACTIVE INGREDIENTS:**

Malathion - O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate . . . . . 5.0%  
Sulfur. . . . . 27.9%

**INERT INGREDIENTS.** . . . . . 67.1%

\*Cythion is a Registered Trade-mark of American Cyanamid Company for the active ingredient Premium Grade Malathion.

## **CAUTION - KEEP OUT OF REACH OF CHILDREN**

Harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin. Skin contact may be harmful. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE TO PHYSICIAN: Malathion is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**DISPOSAL WARNING:** Do not reuse this container. Completely empty the contents. Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

**OBSERVE USE PRECAUTIONS:** Observe use limitations given on the label for specific crops. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

**DIRECTIONS FOR USE:** Read the entire label and conditions of sale, including Warranty Notice, before using Malathion Sulfur. Apply when insects first appear and repeat at 7-10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. **ALFALFA:** Aphids, Use 30 lbs. per acre. No time limitations. **BEANS:** Mexican Bean Beetle, Use 30 lbs. per acre. Do not apply within 1 day of harvest. **CABBAGE-LETTUCE:** Aphids, Leafhoppers, Cabbage Looper. Use 30 lbs. per acre. Do not use within 7 days of harvest. **POTATOES:** Aphids, Leafhoppers, False Cinch Bug. Use 30 lbs. per acre. No time limitation.

**WARRANTY NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured by SCHALL CHEMICAL INC.

"EPA Reg. No. 3468-26"

Monte Vista and Delta, Colorado

ACCEPTED WITH COMMENTS