

# Sevin\* Maneb Sulfur

## ACTIVE INGREDIENTS:

*Carbaryl (1-naphthyl N-methylcarbamate) . . . . .	5.9%
Maneb (Manganese ethylenebisdithiocarbamate) . . . . .	1.0%
Sulfur . . . . .	27.1%

## INERT INGREDIENTS:

61.1%  
\*Sevin, U.S. Pat. No. 2,924,753, 3,099,855. Trademark of Union Carbide Corporation for the active ingredient carbaryl.

## Caution KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin. Skin contact may be harmful. Wash hands and face before eating. Take shower or bath after work. Wear long sleeved work clothing. Change to clean clothing daily. NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

IMPORTANT: Keep away from fire or sparks. Never allow Sevin Maneb Sulfur to become wet during storage. This may lead to certain chemical changes which will reduce effectiveness of Maneb as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Use dust mixture as soon as possible after preparation.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

DISPOSAL WARNING: Do not reuse this container. Completely empty the contents. Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DIRECTIONS FOR USE: Read the entire label and conditions of sale, including Warranty Notice, before using Sevin Maneb Sulfur. Apply dust when disease or insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. POTATOES: Colorado Potato Beetle, Flea Beetle, Leafhopper, Fall Armyworms, Stink Bugs, Tarnished Plant Bugs, Early and Late Blight. Use 20-40 lbs. per acre. No time limitations.

WARRANTY NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured by SCHALL CHEMICAL INC.

E.P.A. Reg. No. 3498 - 27

Monte Vista and Delta, Colo.

