

CYTHION* - MANEB - SULFUR

ACTIVE INGREDIENTS:

Malathion - 0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate. 5.0%
Maneb (manganese ethylenebisdithiocarbamate). 6.0%
Sulfur. 27.9%

INERT INGREDIENTS:

Cythion is a Registered Trade-mark of American Cyanamid Co. for the active ingredient Premium Grade Malathion. 61.1%

CAUTION: KEEP OUT OF REACH OF CHILDREN

May irritate eyes, nose, throat and skin. May be harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. Skin contact may be harmful. Wash hands and face before eating or smoking. Take shower or bath after work or handling. Wear regular long sleeved work clothing. Change to clean clothing daily. In case of contact, flush skin or eyes with plenty of water, for eyes get medical attention. NOTE TO PHYSICIAN: Cythion (Malathion) is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

IMPORTANT: Keep away from fire or sparks. Never allow Maneb to become wet during storage. This may lead to certain chemical changes which will reduce effectiveness of Maneb as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Use dust mixture as soon as possible after preparation.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment, or disposal of wastes. Apply this product only as specified on this label.

This Product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSAL WARNING: Do not reuse this container. Completely empty the contents. Dispose of wastes by burying in noncrop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

OBSERVE USE PRECAUTIONS: Observe use limitations given on the label for specific crops. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

DIRECTION FOR USE: Read the entire label and conditions of sale, including Warranty Notice, before using Cythion-Maneb-Sulfur. Apply when insects or disease first appear and repeat at 7-10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. **CABBAGE:** Aphids, Leafhoppers, Cabbage Looper, alternaria leaf spot, downy mildew. Use 30 lbs. per acre. Do not use within 7 days of harvest. **LETTUCE (Head):** Aphids, leafhoppers, cabbage looper, downy mildew. Use 30 lbs per acre. Remove residues from head lettuce by stripping and trimming. Do not apply within 10 days of harvest. **POTATOES:** Aphids, leafhoppers, false chinch bug, early and late blight. Use 30 lbs per acre. Begin when plants are 2-6 inches high. Repeat at 5-10 day intervals, more often under severe late blight conditions. No time limitations.

WARRANTY NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured by SCHALL CHEMICAL INC.

EPA REG. NO. 3468-31

Monte Vista and Delta, Colorado

