

PRECAUTIONARY STATEMENT—HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Endosulfan is a central nervous system stimulant. There is no specific antidote. Barbituric acid derivatives may be used for convulsions. Treat symptomatically.

Long Pants, long sleeve shirt, gloves, hat and boots must be worn during mixing or flooding.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in cool, dry location. Do not store in direct, hot sunlight. Keep away from fire or sparks. Never allow Maneb to become wet during storage. This may lead to certain chemical changes which will reduce effectiveness of Maneb as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Use dust mixture as soon as possible after preparation.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**BEST DOCUMENT AVAILABLE
THIODAN MANEB SULFOR**

ACTIVE INGREDIENTS:

Thiodan (Thiodan) 27.9%
Maneb (Maneb) 6.0%
Sulfur (Sulfur) 27.9%
Manganese equivalent as metallic 1.23%

INERT INGREDIENTS:

63.1%

* Thiodan is a registered trademark of Canadian Hoechst Limited.
Thiodan U.S. Pat. No. 2,799,685

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed, induce vomiting with 1-2 cups of water or 1/2 cup of 3% hydrogen peroxide. If on skin, wash immediately with soap and water. Consult a physician immediately.

See Side Panel for additional Precautionary Statements.

EPA Reg. No. 3445-38 EPA Est. No. 1068 CO 1
ACCEPTED
 Manufactured by **SCHALL CHEMICAL, INC.**
 Monte Vista, Colorado 81164
SEP 23 1983
 U.S. Dept. of Agriculture
 Agricultural Research Service
 National Plant Disease Survey
 1011 N. 1st St., Fargo, ND 58102
 701/775-3483

DIRECTIONS FOR USE

Apply this dust mixture per acre when necessary to control the above listed insects and diseases. Repeat applications as needed. Observe the following use patterns. Insect control is required beyond the following use patterns. Supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following crop. **CABBAGE, CAULIFLOWER:** (14) Alternaria leaf spot, Downy mildew, Cabbage Aphid, Cabbage Looper, Flea beetles, Harlequin Bug, Sink Bugs, Diamondback Moth Larva, 25-33 lbs per acre. **LETTUCE (Head):** (14) Downy Mildew, Cabbage Looper, Diamondback Moth Larva, Green Peach Aphid, 30-33 lbs per acre. Do not exceed 3 applications on Head lettuce after thinning. Remove outside leaves at harvest. Do not feed crop waste to livestock. **MELONS:** (5) Downy Mildew, Anthracnose, Gummy stem blight, Green Peach Aphid, Cucumber Beetle, 25-33 lbs per acre. Cabbage Looper, 35-50 lbs per acre. Repeat at 7-10 day intervals. **POTATOES:** (0) Early and Late Blight, Aphids, Armyworms, Green Stink Bugs, Plant Bugs, 30-35 lbs per acre. False Chinch Bug, Potato Psyllid, 35 lbs per acre. Repeat at 7-10 day intervals. Do not plant root crops other than carrots, potatoes and sweet potatoes as follow-up crops. **SPINACH:** (21) Downy Mildew (blue mold), Green Peach Aphid, 25 lbs per acre. Do not exceed one application per season. **TOMATOES (Field and Greenhouse):** (5) Early and late blight, Blister Beetles, Colorado Potato Beetle, Flea Beetles, Green Peach Aphid, 20 lbs per acre. Cabbage Looper, 25 lbs per acre. Sink Bug, 20-33 lbs per acre. Tomato Fruitworm, Yellow Striped Armyworm, 35 lbs per acre. Tomato Hornworm, 20-33 lbs per acre. To avoid damage, do not use on tender young greenhouse plants.

NOTE: This pesticide is toxic to fish and other aquatic life. Do not apply to water.

NOTICE — READ CAREFULLY LIMITED WARRANTY STATEMENT

Schall Chemical, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Schall Chemical, Inc. neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. In the event of any breach by Schall Chemical, Inc. of this limited warranty, Schall Chemical, Inc. shall not be liable for consequential damages.

Colson Printing Co. 14556