PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food.

Endangerment [illegible]

[illegible]

PERSONAL PROTECTIVE EQUIPMENT

[illegible]

ENVIRONMENTAL HAZARD

[illegible]

THIODAN MANEB SULFUR

ACTIVE INGREDIENTS:

[illegible] 3.0%

[illegible] 6.0%

[illegible] 27.9%

INERT INGREDIENTS 63.1%

[illegible] Corporation [illegible]

[illegible] 27 [illegible]

MAY [illegible]

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

[illegible] water. Call [illegible]

[illegible]

EPA Reg. No. [illegible]

[illegible] CHEMICAL [illegible]

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in cool, dry location. Do not store in direct, hot sunlight. Keep away from fire or sparks. Never allow Maneb to become wet during storage. This may lead to certain chemical change which will reduce the effectiveness of Maneb as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Use dust mixture as soon as possible after preparation.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of the Federal law to use this product in a manner inconsistent with its labeling. Do not use this product at higher than recommended rate.

[illegible] apply this product in [illegible] manner as to directly [illegible] expose workers or other persons. The [illegible] treated must be [illegible] by unprotected [illegible]

[illegible] is a dust which should be applied in the form with standard dust [illegible] equipment, at the [illegible] each crop. Use [illegible] are for light [illegible] pest and disease pressure. [illegible] the high [illegible] insect and disease pressure.

[illegible] susceptible to sulfur under certain [illegible] conditions. Do not use sulfur on any crop [illegible] unless it has proved that sulfur is safe in that [illegible] high temperatures sulfur may [illegible] Do not apply when temperatures [illegible] exceed 90°F. Do not apply sulfur with [illegible] within 2 weeks of an oil spray treatment. When [illegible] for processing [illegible] processor [illegible] sulfur.

Apply [illegible] first appearance [illegible] is required to [illegible] leaf surfaces is essential for good control. Observe use limitations. If insects or disease control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in ( ) following crop. CABBAGE, (14) Alternaria leaf spot, Downy Mildew, Cabbage Aphid, Cabbage Looper, Flea Beetles, Harlequin Bug, Stink Bugs, Diamondback Moth Larva, 25-33 lbs. per acre. LETTUCE (Head), (14) Downy Mildew, Cabbage Looper, Diamondback Moth larva, Green Peach Aphid, 30-33 lbs. per acre. Do not exceed 3 applications on Head Lettuce after thinning. Remove residues from head lettuce by stripping and trimming outside leaves at harvest. Do not feed crop waste to livestock. MELONS (5) Downy Mildew, Anthracnose, Gummy stem blight, Green Peach Aphid, Cucumber Beetle 25-33 lbs. per acre. Cabbage looper 35-50 lbs. per acre. Repeat at 7-10 day intervals. POTATOES (0) Early and late blight, Aphids, Army worms, Green Stink Bugs, Plant Bugs 30-35 lbs. per acre. False Chinch Bug, Potato Psyllid 35 lbs. per acre. Repeat at 7-10 day intervals. Do not plant root crops other than carrots, potatoes and sweet potatoes as follow up crops. SPINACH (21) Downy Mildew (blue mold), Green Peach Aphid, 25 lbs. per acre. Do not exceed one application per season. TOMATOES (Field and Greenhouse) (5) Early and late blight, Blister Beetles, Colorado Potato Beetle, Flea Beetles, Green Peach Aphid, 20 lbs. per acre. Cabbage Looper 25 [illegible] per acre. Stink Bug 20-33 lbs. per acre. Tomato Fruitworm, Yellow Striped Army worm, 35 lbs. per acre. Tomato Hornworm 20-33 lbs. per acre. To avoid damage, [illegible] not use on tender young growth or seedlings.

NOTE: This [illegible] is to be sold ONLY in the [illegible] unbroken [illegible]

NOTICE - READ CAREFULLY
LIMITED WARRANTY STATEMENT

Schall Chemical [illegible] warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions and instructions on the label. Schall Chemical [illegible] neither makes any other warranty or guarantee or has authorized any agent or representative to make any other warranty of FITNESS, MERCHANTABILITY, or guarantee or representation [illegible] regarding this material. In the event [illegible] by Schall Chemical [illegible] Schall not be liable for consequential damages.

BEST AVAILABLE COPY