#### **OBSERVE THE FOLLOWING USE PRECAUTIONS**

Observe use limitations given on the label for specific crops. BHC may cause off-flavor and odor to certain edible crops.

TO PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.


Apply in early morning or evening when there is little or no

This product is toxic to fish, birds, and other wildlife. Keep out of takes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of smoke).

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing, will vary with local conditions.





(Contains 1 lb. Gamma BHC per Gal'on) INGREDIENTS

**ACTIVE:** Benzene hexachloride, gamma isomer..... 11.83% Benzene hexachloride, other isomers ..... 14.47% Aromatic petroleum derivative solvent . . . . . . 68.70% 

# Keep Out of Reach of Children


May be fatal if swallowed. May produce irritation of skin or eves. May be absorbed through skin! Do not breathe spray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soop and water; for eyes, flush with water and get medical attention. Avoid contamination of food, feed, seed, and

Do Not Store Near Any Product Which Will Absorb Odors! **WARNING. FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME** 

> Protect From Temperatures Below 32° F. **NET CONTENTS**

Manufactured by SCHALL CHEMICAL, INC. Monte Vista, Colorado

Delta, Colorado



## DIRECTIONS FOR USE

Pour specified amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

## RECOMMENDATIONS

APPLES: Aphids — Use 1 qt. in 100 gals. Apply before 60 days of harvest or not after fruit starts to form.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

#### **OBSERVE THE FOLLOWING USE PRECAUTIONS**

Observe use limitations given on the label for specific crops. BHC may cause off-flavor and odor to certain adible crops.

TO PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

Apply in early morning or evening when there is little or no


This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of smoke).

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing, will vary with local conditions.





# DIRECTIONS FOR USE

Pour specified amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

## RECOMMENDATIONS

APPLES: Aphids — Use 1 qt. in 100 gals. Apply before 60 days of harvest or not after fruit starts to form.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

ACCEPTED 467-39 OCT 31, 1973

UNDER THE "SPERAL IN" FUNGICIDI. FOR ECONUP 1 EQ UNDER N' TO ATTACHED TO