

CAUTION Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water, for eyes get medical attention. Keep out of reach of children. Harmful if swallowed.

CONTAINER DISPOSAL. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

CAUTIONS. Do not contaminate water by cleaning of equipment, or disposal of wastes, and apply this product only as specified on this label.

IMPORTANT. Prevent drift of product to desirable plants, particularly soybeans (Coarse sprays are less likely to drift). For airplane application use 3 or more gallons of water per acre. Do not apply by airplane when sensitive crops are nearby. Consult your state or local authorities for possible aerial application or other application restrictions. For ground application use sufficient water to give good coverage. Do not store near fertilizers, seeds, insecticides, or fungicides. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not use same spray equipment for other purposes. Avoid spray drift to susceptible plants, as this product may injure cotton, beans, peas, grapes, ornamentals, etc. Do not contaminate water used for irrigation or domestic purposes. Do not apply (except as recommended) on or near desirable trees or plants, or in locations where chemicals may be washed or moved into contact with their roots.


USE DIRECTIONS
To minimize BANTROL spray drift onto potatoes and other sensitive crops, the following suggestions are given.

1. Use coarse sprays by decreasing the spray pressure below 20 p.s.i.
2. Spray with the nozzles as close to the weeds as is practical for good coverage.
3. Do not spray when wind is over 5 miles per hour.
4. If any wind is blowing in the direction of the potato field, do not spray.

For ground equipment use 10 to 20 gallons of water per acre when treating broadleaf weeds and for tip growth control of perennial broadleaf weeds. For medium control of perennial broadleaf weeds, use 100 gallons or more of water per acre.

BANTROL

EPA Reg. No. 3468-47
EPA EST: 3468-CQ-1



ACTIVE INGREDIENTS:

- * Dimethylamine Salt of dicamba (3, 6-dichloro-o-anisic acid) 12.25%
- Dimethylamine Salt of related acids 1.97%
- ** Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic Acid . . . 24.40%
- * Equivalent to 10.15% 3, 6-dichloro-o-anisic acid.
- ** Equivalent to 20.3% 2, 4-dichlorophenoxyacetic acid. **3468-47**
- INERT INGREDIENTS 61.38%**

Net Contents _____ Gallons

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel For Additional Cautions.

Manufactured by SCHALL CHEMICAL, INC.

Monte Vista, Colorado

BARLEY, OATS WHEAT (Fall) for the control of Canada Thistle, Pigweed, Ragweed, Smartweed, 1 pint per acre. Apply as a foliar dormancy and before the grain is actively growing. Do not graze prior to crop maturity. Do not graze in Arizona.

PASTURE and RANGELAND of fence rows, roadways and waste wheat, Annual clover, Lamb's ear, ragweed, Rainbowbrush and Sunflower, dock, Kochia, Morningglory (sp), use 2 quarts of Bantrol per acre. **Russian Thistle Field bindweed, Leafy spurge and Perennial sow** Wild garlic, Goldenrod, Poverty are actively growing. Do not graze. Do not apply to newly seeded. For ground equipment use 10-20 annual broadleaf weeds and for leaf weeds. For maximum control use 100 gallons or more of water per acre. Oklahoma and New Mexico. Do not graze pastures and clover and

Do not graze meat animals with. Do not graze dairy animals on. Do not apply within 30 days of the seed from treated grass.

NOTICE: Seller's guarantee and subject there to the buyer arising out of use or handling.


BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

case of contact,
 keep out of
 sh thoroughly
 ent, or dia-
 this label.
 particularly
 ne application
 plane when
 orities for
 For ground
 at store near
 lean spray
 other purposes,
 id spray drift
 us, peas, grapes,
 m or domestic
 desirable
 ad or moved
 ensitive crops,
 w 20 p.e.l.
 val for good
 do not spray.
 when treat-
 roadleaf
 use 100 gal-

BANTROL

EPA Reg. No. 3468-47
 EPA EST: 3468-CQ:1



ACTIVE INGREDIENTS:

* Dimethylamine Salt of dicamba (3, 6-dichloro-o-anisic acid)	12.25%
Dimethylamine Salt of related acids	1.97%
** Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic Acid	24.40%
* Equivalent to 10.15% 3, 6-dichloro-o-anisic acid.	
** Equivalent to 20.3% 2, 4-dichlorophenoxyacetic acid.	
INERT INGREDIENTS	61.38%

Net Contents _____ Gallons

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel For Additional Cautions.

Manufactured by SCHALL CHEMICAL, INC.

Monte Vista, Colorado

BARLEY, OATS WHEAT (Fall seeded) (not underseeded to legumes) for the control of Canada Thistle (seedling or top growth) Lambsquarter, Pigweed, Ragweed, Smartweed, Sowthistle (annual) Wild buckwheat use 1 pint per acre. Apply as a foliar spray immediately after winter dormancy and before the grain begins to joint. Apply when weeds are actively growing. Do not graze treated areas or harvest for dairy feed prior to crop maturity. Do not use on Fall Seeded Barley in California or Arizona.

PASTURE and RANGELAND GRASSES and NONCROPLANDS, such as fencerows, roadways and wastelands. For the control of Wild buckwheat, Annual clover, Lambsquarters, Mustard, Redroot pigweed, Common ragweed, Rattlebrush and Sunflower use 2 pints Bantrol per acre. Curly dork, Kochia, Morningglory (annual) Puncturevine and Shepherdspurse use 2 quarts of Bantrol per acre. Top Growth Control of Canada Thistle, Russian Thistle Field bindweed, Black knapweed, Russian knapweed, Leafy spurge and Perennial sowthistle use 2 quarts per acre. Clover, Wild garlic, Goldenrod, Poverty weed and Sowthistle use 1 gallon per acre. For control or suppression of listed weeds, apply Bantrol when weeds are actively growing. Do not apply when grass is in boot to milk stage. Do not apply to newly seeded areas after heading begins. For ground equipment use 10-20 gallons of water per acre when treating annual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. For Use Only in the States of Texas, Oklahoma and New Mexico. Do not use this mixture on Bent, susceptible grass pastures and clover and alfalfa pastures.

Do not graze meat animals within 30 days of slaughter.
 Do not graze dairy animals on treated areas with 42 days of slaughter.
 Do not apply within 70 days of harvest.
 The seed from treated grass should not be used for feed or food purposes.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

IT AVAILABLE

BEST DOCUMENT AVAILABLE

CAUTION Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Keep out of reach of children. Harmful if swallowed.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

CAUTIONS: Do not contaminate water by cleaning of equipment, or disposal of wastes, and apply this product only as specified on this label.

IMPORTANT: Prevent drift of product to desirable plants, particularly soybeans. (Coarse sprays are less likely to drift). For airplane application use 3 or more gallons of water per acre. Do not apply by airplane when sensitive crops are nearby. Consult your state or local authorities for laws for aerial application or other application restrictions. For ground equipment use sufficient water to give good coverage. Do not store near fertilizers, seeds, insecticides, or fungicides. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes, or do not use same spray equipment for other purposes. Avoid spray drift to susceptible plants, as this product may injure cotton, beans, peas, grapes, ornamentals, etc. Do not contaminate water used for irrigation or domestic purposes. Do not apply, except as recommended, on or near desirable trees or plants, or in locations where chemicals may be washed or moved into contact with their roots.

USE DIRECTIONS
To minimize BANTROL spray drift onto potatoes and other sensitive crops, the following suggestions are given:

1. Use coarse sprays by decreasing the spray pressure below 20 p.s.i.
2. Spray with the nozzles as close to the weeds as is practical for good coverage.
3. Do not spray when wind is over 5 miles per hour.
4. If any wind is blowing in the direction of the potato field, do not spray.

For ground equipment use 10 to 20 gallons of water per acre when treating broadleaf weeds and for top growth control of perennial broadleaf weeds. For airplane control of perennial broadleaf weeds, use 100 gallons or more per acre.

BANTROL

EPA Reg. No. 3468-47
EPA EST. 3468-CQ-1

ACTIVE INGREDIENTS:

* Dimethylamine Salt of dicamba (3, 6-dichloro-o-anisic acid)	12.25%
Dimethylamine Salt of related acids	1.97%
** Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic Acid	24.40%
* Equivalent to 10.15% 3, 6-dichloro-o-anisic acid.	
** Equivalent to 20.3% 2, 4-dichlorophenoxyacetic acid.	
INERT INGREDIENTS	61.38%

Net Contents _____ Gallons

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel For Additional Cautions.

Manufactured by SCHALL CHEMICAL, INC.

Monte Vista, Colorado


BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVA

BAR...
for...
Fig...
I pr...
dom...
acti...
DIME...
Ariz...
PAST...
fenc...
rag...
dore...
use...
Rus...
Le...
Wid...
For...
Are...
Do...
For...
and...
LOW...
ON...
gra...

BANTROL

EPA Reg. No. 3468-47
EPA EST. 3468-CQ-1



ACTIVE INGREDIENTS:

Dimethylamine Salt of dicamba (3, 6-dichloro-o-anisic acid)	12.25%
Dimethylamine Salt of related acids	1.97%
* Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic Acid	24.40%
* Equivalent to 19.15% 3, 6-dichloro-o-anisic acid.	
** Equivalent to 20.3% 2, 4-dichlorophenoxyacetic acid.	
INERT INGREDIENTS	61.38%

Net Contents _____ Gallons

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel For Additional Cautions.

Manufactured by SCHALL CHEMICAL, INC.

Monte Vista, Colorado

BARLEY, OATS WHEAT (Fall seeded) (not underseeded to legumes) for the control of Canada Thistle (seedling or top growth) Lamb's-quarter, Pigweed, Ragweed, Smartweed, Sooty thistle (annual), Wild buckwheat use 1 pint per acre. Apply as a foliar spray immediately after winter dormancy and before the grain begins to joint. Apply when weeds are actively growing. Do not graze treated areas or harvest for dairy feed prior to crop maturity. Do not use on Fall Seeded Barley in California or Arizona.

PASTURE and RANGELAND GRASSES and NONCROPLANDS, such as fencerows, roadways and wastelands. For the control of Wild buckwheat, Annual clover, Lamb's-quarters, Mustard, Red top pigweed, Common ragweed, Ragwort, and Sunflower use 2 quarts Bantrol per acre. Curry dock, Knapweed, Morning glory (annual), Purple spurge and Shepherd's-purse use 2 quarts of Bantrol per acre. Top Growth Control of Canada Thistle, Russian Thistle, Field bindweed, Black knapweed, Russian ragweed, Leafy spurge and Perennial sowthistle use 2 quarts per acre. Clover, Wild garlic, Goldenrod, Poverty weed and Sowthistle use 1 gallon per acre. For control or suppression of listed weeds, apply Bantrol when weeds are actively growing. Do not apply when grass is in seed or milk stage. Do not apply to newly seeded areas after seeding begins. For ground equipment use 10-20 gallons of water per acre when treating annual broadleaf weeds and for top growth control of perennial broadleaf weeds. For maximum control of perennial broadleaf weeds use 100 gallons or more of water per acre. For Use Only in the States of Texas, Oklahoma and New Mexico. Do not use this mixture on bent, susceptible grass pastures and clover and alfalfa pastures.

Do not graze meat animals within 30 days of slaughter.
Do not graze dairy animals on treated areas within 42 days of slaughter.
Do not apply within 70 days of harvest.
The seed from treated grass should not be used for feed or food purposes.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

BEST DOCUMENT AVAILABLE