


(Crop Dust)

# ZINEB - SULFUR



**ACTIVE INGREDIENTS:**

Zineb (zinc ethylene bisdithiocarbamate) . . . . .	7.0%
Sulfur . . . . .	50.0%

**INERT INGREDIENTS . . . . .**

43.0%

## CAUTION: KEEP OUT OF REACH OF CHILDREN

May irritate eyes, nose, throat and skin. May be harmful if inhaled or swallowed. Avoid breathing of dust. ~~Avoid contact with skin, eyes or clothing.~~ Skin contact may be harmful. Wash hands and face before eating or smoking. Take shower or bath after work or handling. Wear regular long sleeved work clothing. Change to clean clothing daily. In case of contact, flush skin or eyes with plenty of water, for eyes get medical attention.

**IMPORTANT:** Keep away from fire or sparks. Never allow Zineb to become wet during storage. This may lead to certain chemical changes which will reduce effectiveness of Zineb as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Use dust mixture as soon as possible after preparation.

**DISPOSAL WARNING:** Do not reuse this container. Completely empty the contents. Dispose of wastes by burying in noncrop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

**ENVIRONMENTAL CAUTIONS:** Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**OBSERVE USE PRECAUTIONS:** Read the entire label and conditions of sale, including Warranty Notice, before using Zineb-Sulfur. Observe use limitations given on the label for specific crops. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others. Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label's cautions and directions.

**DIRECTIONS FOR USE:** Apply when disease first appears and repeat at 7-10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. **LETTUCE:** (Head) apply with ground dusting equipment only and only when both downy mildew and powdery mildew are expected to be controlled. Apply 30 lbs. per acre when both these diseases first appear and repeat at 7-10 day intervals as required to maintain control. Do not use within 10 days of harvest. Sulfur may burn foliage when temperature is high. Do not make sulfur applications at such times.

**WARRANTY NOTICE:** Seller's guarantee shall be limited to the terms of the label, and subject there to the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.