

ACCEPTED
DEC 9 1966
UNDER THE FEDERAL INSECTICIDE
REGULATION ACT REGISTERED
FOR ELEM. NO. 2487-19 SUBJECT
ED UNDER INVOICE NO. 2487-19
TO ATTACHED COMMENTS



Place tablespoon amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8-12 feet are recommended. Assure uninterrupted supply of bait for a period of not less than 12 days.

Place 4 to 6 ounces of bait in dry containers next to burrows, in runways or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. **OPEN SAUCERS OF BAIT IS PROPER ONLY WHEN THE AREA IS INACCESSIBLE TO CHILDREN, PETS, AND DOMESTIC ANIMALS. OTHERWISE, THE BAITS SHOULD BE EXPOSED UNDER PROTECTIVE COVER THAT ONLY A RAT OR MOUSE CAN ENTER.**

Replace stale, damp and moldy baits with fresh baits.

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood, and upon repeated feedings, causes fatal hemorrhage in rats. It also contains an antibacterial agent, sulfaquinoxaline, which inhibits bacterial that produce Vitamin K, an antidote for warfarin. It thereby increases effectiveness against hard-to-kill rats and mice and gives better more thorough control.

MSD # 3487-X



BACON PRODUCTS, INC.
P. O. BOX 8127
CHATTANOOGA, TENNESSEE

CAUTION Keep out of reach of Children
See back panel for additional cautions

Place tablespoon amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8-12 feet are recommended. Assure uninterrupted supply of bait for a period of not less than 12 days.

Place 4 to 6 ounces of bait in dry containers next to burrows, in runways or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. **OPEN SAUCERS OF BAIT IS PROPER ONLY WHEN THE AREA IS INACCESSIBLE TO CHILDREN, PETS, AND DOMESTIC ANIMALS. OTHERWISE, THE BAITS SHOULD BE EXPOSED UNDER PROTECTIVE COVER THAT ONLY A RAT OR MOUSE CAN ENTER.**

Replace stale, damp and moldy baits with fresh baits.

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood, and upon repeated feedings, causes fatal hemorrhage in rats. It also contains an antibacterial agent, sulfaquinolone, which inhibits bacterial that produce Vitamin K, an antidote for warfarin, and thereby increases effectiveness against hard-to-kill rats and mice and gives better more thorough control.



Net Weight 1 lb.

98¢

Eagle's 7

RAT BAIT

**KILLS
RATS & MICE**

CONTAINS

U. S. Patent Nos. 2,487,578 and 2,111,871
A better, more effective warfarin rodenticide

ACTIVE INGREDIENTS:	
9-(4-oxo-1,2,3,4-tetrahydroquinolin-2-yl)-4-hydroxycoumarin (warfarin)	0.025%
N1-2-quinazolinylsulfanilamide (sulfaquinolone)	0.025%
INERT INGREDIENTS	99.950%
	100.000%



BACON PRODUCTS, INC.

P. O. BOX # 27

WALTONOOGA, TENNESSEE