

3487-19

3/24/2004

DIRECTIONS FOR USE

1/1

Eagles-7 Rat Bait

Kills Norway & Roof Rats House Mice

ACTIVE INGREDIENT:

Warfarin..... 0.025%

OTHER INGREDIENTS..... 99.975%

Total..... 100.000%

EPA Reg. No. 3487-19

EPA Est. No. 2497-FA

Net Wt. 4 oz (113g)

Net Wt. 6 oz (170g)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautionary Statements

ACCEPTED
 MAR 24 2004
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide WARFARIN under EPA Reg. No. 3487-19

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized.
3. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
4. Dispose of product container and unused, spoiled and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice in and around homes, industrial buildings and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Apply 4 to 16 oz. of bait per placement. This package contains enough bait for 2 or 2 rats. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE: Apply 1/4-1/2 oz. (1-2 tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 oz.) may be needed at points of extremely high mouse activity. Space placements at intervals of 8-12 feet. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Collect and properly dispose of all dead animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Wear gloves if you handle bait.

Environmental Hazards

This product is toxic to mammals and birds. Do not apply directly to water.

FIRST AID

Have label with you when obtaining treatment advice.

If swallowed

- Call a poison control center, doctor, or 1-800-222-1222 immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes
- Call a poison control center, doctor, or 1-800-222-1222 immediately for treatment advice.

NOTE TO PHYSICIAN

Warfarin may reduce the clotting ability of the blood and cause bleeding. If ingested, Vitamin K₁ may need to be given orally or intramuscularly. Repeat as needed, based on prothrombin times.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured by: Bacon Products Corporation
P O Box 22187
Chattanooga, TN 37422