

## OFFICE OF PESTICIDE REGISTRATION

(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

## NAME OF PESTICIDE PRODUCT

Eagles-7 Lawn and Garden Insect  
Control

## NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Bacon Product Corporation  
P.O. Box 22187  
Chattanooga, TN 37422-2187160 / 145439  
18 / 1

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 3487-75" to your label before you release the product for shipment.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. The additional brand name, "Fire Fighter" is also being accepted at this time.

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

Enclosure

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL



DATE

3/24/75

MAR 28 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

2487-25

# Eagle-7

## lawn & garden insect control

# KILLS

Ants  
Armyworms  
Bermudagrass  
mites  
Billbugs  
Brown dog  
ticks  
Chiggers  
Chinch bugs  
Crickets  
Cutworms  
Earwigs

Ready-to-use granules.  
Contains 5% diazinon  
insecticide.

Keep Out of Reach  
of Children.

### Caution

See additional precautionary state-  
ments on back of bag.

Active ingredient:  
Diazinon: O, O-diethyl O-(2-  
isopropyl-8-methyl-4-pyrimid-  
inyl) phosphorothioate ..... 5%  
Inert ingredients: ..... 95%  
Total: 100%

10 lbs.  
Net Weight

Fire Ants\*  
Fleas  
Onion maggots  
Root maggots  
Sod webworms  
White grubs of:  
• Japanese beetle  
• European chafer  
• Southern chafer  
Wireworms  
and others

\*Aids in control.  
See Directions for Use.

**In Lawns**

*Eagle-T* kills:

- Ants
- Armyworms
- Bermudagrass mites
- Billbugs
- Brown dog ticks
- Chiggers
- Chinch bugs
- Clover mites
- Crickets
- Cutworms
- Digger wasps
- Earwigs
- Fire Ants\*
- Fleas
- Lawn moths
- Leafhoppers
- Mistlebees
- Sod webworms
- Sowbugs
- Springtails
- White grubs of:
  - Japanese beetle
  - European chafer
  - Southern chafer

\*Aids in control.

**In Gardens**

*Eagle-T* kills:

- Armyworms
- Carrot rust flies
- Corn borers
- Cutworms
- Mole crickets
- Onion maggots
- Root maggots
- Rootworms
- Vinegar flies
- Wireworms

It is a violation of Federal law to use this product on a residential lawn except as labeled.

**Overall Lawn Application:**  
Apply Eagle-T through the following lawn insects on both grass and clover lawns:  
Ants - Armyworms - Bermudagrass mites - Brown dog ticks - Chiggers - Chinch bugs - Clover mites - Crickets - Cutworms - Earwigs - Fleas - Lawn moths (and cutworms) - Leafhoppers - Mistlebees - Sowbugs - Springtails (Collembola)

Apply Eagle-T through a fertilizer spreader. See spreader setting chart on side of bag. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs per 1000 sq ft lawn area by 20" or 3.2 oz per 100 sq ft lawn area by 10". For heavy lawns such as St Augustine grass lawns, the rate should be increased to 2 lbs per 1000 sq ft (or 4.8 oz per 100 sq ft).

**Billbugs:** White grubs of Japanese beetle, European chafer, Southern chafer, Atlantic ground (living) beetle:  
Apply 2.5 lbs per 1000 sq ft or 1/2 cup Water granules through a fertilizer spreader. For billbug control, apply when activity is first observed or when showed in brown grass indicates damage from these insects. Repeat as necessary. For grub control, control June beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water granules thoroughly with 1/2 in of water after application. Treat again in late summer or early fall if a second generation occurs.

**Hyperodes weevils (New York only):**  
Apply 2 lbs per 1000 sq ft of grass. Water granules thoroughly after application. This product active in mid-April and again in mid-May.

**Spot Treatment**  
Ants: Apply 1/2 oz of Eagle-T around anthills over and around each anthill.

**Fire ants:** To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over and 1/2 in around each fire ant mound immediately after application. Water thoroughly. A minimum of 1 gallon of water should be applied to each mound. Repeat as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment will disturb the ants and cause migration (reduce) product effectiveness.

**Note:** For best results, apply in cool weather, 60-80 F, in an early morning or late evening hours. Treat new mounds as they appear.

**Digger wasps:** Apply 1 1/2 teaspoons of Eagle-T in a 3-foot band in and around each nest, covering. Apply a dust when wasps are not active.

**Band Treatment Around House Foundation:**  
The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses:

- | Ants            | Fleas              |
|-----------------|--------------------|
| Brown dog ticks | Mistlebees         |
| Clover mites    | Sowbugs (Pillbugs) |
| Crickets        | Springtails        |
| Sowbugs         | (Collembola)       |

Apply a 3-foot band of Eagle-T around to the house foundation. Distribute granules uniformly using a rate of 3.2 oz for each 100 sq ft of area (5" by 20") treated. Repeat as necessary.

**Garden Insect Control:**  
Apply Eagle-T through a fertilizer spreader on a vegetable garden on a regular basis as labeled below. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth, except where applied in seed furrows.

- | Vegetables (surface and subterranean) | Other      | Other         |
|---------------------------------------|------------|---------------|
| Broccoli                              | Kale       | Snap Beans    |
| Cabbage                               | Letuce     | Spruce        |
| Cantaloupe                            | Lima Beans | Summer Squash |
| Cauliflower                           | Muskmelons | Sweet Chard   |
| Celery                                | Peas       | Tomatoes      |
| Collards                              | Peppers    | Watermelon    |
| Cucumbers                             | Potatoes   | Winter Squash |
| Endive                                | (Escarole) |               |

Apply 7 to 14 oz per 500 sq ft and mix into soil to a depth of 2-3 inches for surface cutworms and 3-4 inches for subterranean cutworms. For potatoes, mix into soil 4-6 inches deep.

- | Other         | Other      | Other        |
|---------------|------------|--------------|
| Mole Crickets | Endive     | Peppers      |
| Broccoli      | (Escarole) | Radishes     |
| Cabbage       | Kale       | Straightneck |
| Cauliflower   | Letuce     | Turnips      |
| Collards      | Peas       |              |

Apply 4 oz per 500 sq ft and mix into soil to a depth of 1-2 inches.

- | Other       | Other      | Other          |
|-------------|------------|----------------|
| Wireworms   | Endive     | Peppers        |
| Broccoli    | (Escarole) | Snap Beans     |
| Cabbage     | Kale       | Spruce         |
| Cantaloupe  | Letuce     | Summer Squash  |
| Carrots     | Lima Beans | Sweet Potatoes |
| Cauliflower | Muskmelons | Sweet Chard    |
| Celery      | Onions     | Tomatoes       |
| Collards    | Peas       | Watermelon     |
| Corn        |            | Winter Squash  |

Apply 7 to 14 oz per 500 sq ft and mix into soil to a depth of 4-8 inches.

**Broccoli, Cabbage, Cauliflower:**  
Root Maggots: Apply 7 to 11 oz per 500 sq ft over entire seedbed before planting or transplanting and mix into soil to a depth of 3-4 inches.

**Carrots:**  
Carrot Rust Flees: Apply 7 oz per 500 sq ft in seed furrow at planting time.

**Onions:**  
Onion Maggots: Apply 4 oz per 500 sq ft in seed furrow at planting time.

**Radishes:**  
Root Maggots: Sprinkle lightly at the rate of 3-4 oz per 500 linear feet of row in seed furrow at planting time.

**Sweet Corn:**  
Corn Rootworms: Apply 4.7 oz per 500 sq ft in a band over the row at planting time.

**Cucumbers:** Apply 7.11 oz per 500 sq ft. Mix into soil to a depth of 1-2 inches for surface cutworms and 3-4 inches for subterranean cutworms.

**Eurobean Corn Beans:** Apply 4.7 oz per 500 sq ft over corn so granules fall into whorls when 75% of the plants show first-generation larvae feeding, and again 7 days later if needed. Treat for second generation when there is an average of 1 egg mass per plant.

**Fat Armyworms:** Apply 4.7 oz per 500 sq ft over corn so granules fall into whorls when 75% of the plants show first-generation larvae feeding, and again 7 days later if needed. Treat for second generation when there is an average of 1 egg mass per plant.

**Spotted-stem Corn Beets:** Apply 4-7 oz per 500 sq ft over corn so granules fall into whorls when second or third instars are first seen. Repeat application only when active.

\*Note: Corn may be picked immediately following last application. Do not feed fodder to livestock for ten days following last application.

**Wireworms:** Apply 7-15 oz per 500 sq ft in a broadcast application. Mix into soil to a depth of 4-6 inches.

**Sweet Potatoes (North Carolina only):**  
Wireworms: Apply 11 oz per 500 sq ft over the top of sweet potato plant foliage when roots begin to enlarge.

**Tomatoes:**  
Vinegar Flies (Drosophila spp.): Apply 4 oz per 500 sq ft over top of plants when first appear. Repeat at 7-10 day intervals, or as required. Tomatoes may be picked the same day.

**Turnips:**  
Root Maggots: Apply 4 oz per 500 sq ft in seed furrow at planting time.

**Watermelons:** Apply 4 oz per 500 sq ft in seed furrow at planting time.

**Winter Squash:** Apply 4 oz per 500 sq ft in seed furrow at planting time.

**STORAGE AND DISPOSAL:**  
Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container, even container and mix in trash collection.

EPA REG NO. 3487-  
EPA EST NO. 3487-01-1

**BACK LABEL FOR: EAGLE-T LAWN AND INSECT CONTROL EPA 3487-E**

**NOTE:**  
The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary in order to secure correct rate of application.

**SPREADER SETTING NO. (per 1000 sq ft.)**

Spreader Type and Brand	Broadcast Rate		Rate (lb/1000 sq ft)
	1/2 in.	1 in.	
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Carton Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1
Hand Broadcast Rate (lb/1000 sq ft)	7 1/2	15	1

**PRECAUTIONARY STATEMENTS:**  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION: If inhaled or absorbed through skin, avoid breathing dust and contact with eyes, skin, and clothing. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.  
STATEMENT OF PRACTICAL TREATMENT  
If swallowed, call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.  
If inhaled, remove victim to fresh air. Apply respiration if indicated.  
If on skin, remove contaminated clothing and wash affected areas with soap and water.  
If in eyes-Flush with plenty of water. Call a physician immediately.  
NOTE TO PHYSICIAN: If symptoms of cholinergic poisoning occur, administer atropine therapeutically.  
ENVIRONMENTAL HAZARDS  
This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.