

SEP - 8 1993

310/5439182
18

Mr. James Edwards
Bacon Products Corporation
P.O. Box 22187
Chattanooga, TN 37422

Dear Mr. Edwards:

Subject: Amendment - Addition of Sites and Pests
Eagles 7 Flea & Tick Spray
EPA Registration No. 3487-27
Your Application Dated March 23, 1993

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the labeling changes listed below before you release the product for shipment:

- a. Delete the statement "Do not apply to classrooms while in use." Institutional sites are not listed on the label.
- b. On the front panel add the statement "For control of Flies and Lice on Livestock and Premises."

3. Submit five copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

Eagles-7 FLEA & TICK Spray

EPA REG. NO. 3487-27
EPA EST. NO. 3487-TN-1

- For Homeowner Use Both Indoors and Outdoors
- A Ready-To-Use PERMANONE® Stable Emulsion Insecticide
- For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs
- For Control of Ticks which may cause Lyme disease on Premises
- Controls Fleas and Ticks on Dogs for 35 Days
- Aids in Control of Ticks on Dogs for an additional 2 weeks

ACTIVE INGREDIENT: *Permethrin.....0.50%
 INERT INGREDIENTS:99.50%
 Total 100.00%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethoxy)2,2-dimethylcyclopropanecarboxylate.
 Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

PERMANONE - Registered Trademark of Fairfield American Corporation.

Net Contents

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of food and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SHAKE WELL BEFORE USING INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors. Do not apply to classrooms while in use. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

SURFACE SPRAYING

To control *Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Caddisflies*: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

To control *Carpet Beetles*: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control *Bedbugs*: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control *Fleas (adults and larvae) and Ticks, including Ticks which may carry Lyme disease (adults and larvae)*: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

To control *Clothes Moths*: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To kill *Lice and House Eggs*: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ACCEPTED
with COMMENTS
in EPA Letter Dated

SEP - 8 1993

BEST AVAILABLE COPY

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

3487-27

ON LIVESTOCK:

To protect cattle (beef and dairy), goats, sheep, hogs and horses from *Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats*: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control *Stable Flies, Horse Flies and Deer Flies*: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control *Blood-sucking Lice*: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control *Poultry Lice*: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control *Bedbugs and Mites* in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS:

To control *Fleas, Ticks and Lice*: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat as necessary.

NOTE: Not recommended for use on puppies less than four weeks old. **DO NOT USE ON CATS OR KITTENS.** Do not use on old, sick or debilitated animals.

OUTDOOR APPLICATION

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):

Use Eagles-7 Flea & Tick Spray as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*), Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*).

For each use areas listed below, apply until surface is wet (approximately 1gal/75-1000ft).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around and the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 foot wide and adjacent to activity areas; especially where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS:

For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray as well as *Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers*. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English Ivy, eunonymus, fuchsia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against *Whiteflies, Aphids, Spiders and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms*.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

Bacon Products Corporation
P. O. Box 22187
Chattanooga, TN 37422-2187 U.S.A.