

3487-27

( 09/28/2012 )

1/11

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D C 20460-0001



OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

September 28, 2012

Ms Karen M Romito  
Bacon Products Corporation  
P O Box 22187  
Chattanooga, TN 37422

Subject Amended Reregistration Label  
Product Name Eagles-7 Flea & Tick Spray  
EPA Registration Number 3487-27  
EPA Decision Number 413479

Dear Ms Romito,

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the reregistration of the above referenced product in connection with the pyrethrin and piperonyl butoxide RED(s), and has concluded that your submission is acceptable

NOTE This product is not being reregistered under sections 3(c)5 and 4(g) of FIFRA at this time

Please be reminded that 40 CFR Part 156 140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on non-refillable containers. The code may appear either on the label or durably marked on the container itself, and can be added by non-notification per PRN 98-10

Please note that the record for this product currently contains the Confidential Statements of Formulation (CSFs) listed below. Any previously dated CSFs are superseded

- Basic CSF, dated October 11, 2011

A copy of your label stamped "Accepted" is enclosed along with copies of the acute toxicity and product chemistry reviews completed for the subject product. Products shipped

after 12 months from the date of this amendment or the next printing of the label whichever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e)

If you have any questions about this letter, please contact Richard Gebken at (703) 305-6701 or via e-mail at [gebken\\_richard@epa.gov](mailto:gebken_richard@epa.gov) or Jennifer Gaines at (703) 305-5967 or via e-mail at [gaines\\_jennifer@epa.gov](mailto:gaines_jennifer@epa.gov)

Sincerely,




Richard J Gebken  
Product Manager (10)  
Insecticide Branch  
Registration Division (7504P)

*Enclosures    Label stamped "Accepted," dated September 28, 2012  
Acute Toxicity Review, dated July 25, 2011  
Product Chemistry Review, dated November 3, 2011*

**SUB-LABEL A – HOMEOWNER**

**Eagles-7 Flea & Tick Spray**



- *[For Homeowner Use Both Indoors And Outdoors*
- *A Ready-To Use Permanone Stable Emulsion Insecticide*
- *For Control Of Fleas, Brown Dog Ticks And Lice On Premises And On Dogs*
- *For Control Of Ticks Which May Cause Lyme Disease On Premises*
- *Controls Fleas And Ticks On Dogs For 35 Days*
- *Aids in Control of Ticks on Dogs for an additional 2 weeks]*

**ACTIVE INGREDIENT**

Permethrin 0 50%

**OTHER INGREDIENTS**

99 50%

100 00%

Cis/trans ratio min 35% (+) cis and max 65% (+) trans

**EPA REG NO 3487 27**

**EPA EST NO**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back/side panel for additional precautions

**NET CONTENTS**

**READ ENTIRE LABEL BEFORE EACH USE  
PET USE APPLIES TO USE ONLY ON DOGS**

**PRECAUTIONARY STATEMENTS  
Hazards To Humans & Domestic Animals**

**CAUTION**

Wash hands thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet Remove and wash contaminated clothing before reuse

**Environmental Hazards**

This product is extremely toxic to aquatic organisms including fish and invertebrates To protect the environment do not allow pesticide to enter or run off into storm drains drainage ditches gutters or surface waters Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

### Use Restrictions & Precautions

- Except when applying to pets do not enter or allow adults children or pets to enter the treated area until sprays have dried
- Not for broadcast use on indoor surfaces
- Cover all food processing surfaces and utensils during treatment or thoroughly wash before use Cover or remove exposed food/feed
- Do not contaminate food or feedstuffs
- Remove pets and birds and cover fish aquaria before spraying and surface applications
- Do not use on electrical equipment or on surfaces that can be harmed or stained by water
- Application is prohibited directly into sewers or drains or to any area like a gutter where drainage to sewers storm drains water bodies or aquatic habitat can occur Do not allow the product to enter any drain during or after application
- For pet care Do not apply to dogs less than 12 weeks old Consult a veterinarian before applying this product on medicated debilitated aged pregnant or nursing animals Sensitivities may occur after using ANY pesticide product for pets If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water If signs continue consult a veterinarian immediately
- After applications outdoors do not water the treated area to the point of run off
- Do not make outdoor applications during rain
- Not for use in outdoor residential misting systems
- All outdoor applications must be limited to spot or crack and-crevice treatments only except for the following permitted uses
  - 1 Treatment to soil or vegetation around structures
  - 2 Applications to lawns turf and other vegetation
  - 3 Applications to building foundations up to a maximum height of 3 feet
- Other than applications to building foundations all outdoor applications to impervious surfaces such as sidewalks driveways patios porches and structural surfaces (such as windows doors and eaves) are limited to spot and crack and crevice applications only

Ready to use Turn nozzle on trigger sprayer to desired pattern Do not transfer to another container

**SHAKE WELL BEFORE USING**

**INDOOR APPLICATION**

For spot application only Do not use as a space spray indoors

**SPOT AND CRACK & CREVICE APPLICATION**

To control of *Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles* Direct the spray into hiding places cracks and crevices and behind shelves and drawers For *Silverfish*, spray bookcases For *Ants*, spray trails nests and points of entry

TO CONTROL *Carpet Beetles* Spray edges of carpeting and under carpeting and rugs Make localized application to floor and baseboards Spray directly into cracks crevices and infested areas of shelving

TO CONTROL *Bedbugs* Spray mattresses lightly particularly around tufts and seams Take beds apart and spray in all joints Treat baseboards moldings and floors

TO CONTROL *Fleas (adults and larvae)* and *Ticks, including Ticks which may carry Lyme disease (adults and larvae)* Thoroughly spray infested areas pet beds resting quarters nearby cracks and crevices along and behind baseboards moldings window and door frames and entire areas of floor and floor covering Place fresh bedding in animal quarters following treatment Allow to dry thoroughly before allowing pets to contact treated areas

TO CONTROL *Clothes Moths* Remove any infested articles from storage brush thoroughly and air for several hours in sunlight if possible Apply spray liberally to empty chests closets bureaus and other storages directing the spray into cracks joints and crevices After airing infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining

TO KILL *Lice* and *Louse Eggs* Spray in an inconspicuous area to test for possible staining or discoloration Inspect again after drying then proceed to spray entire area to be treated Spray from a distance of 8 to 10 inches Treat only those garments and parts of bedding including mattresses and furniture that cannot be either laundered or dry cleaned Allow all treated articles to dry thoroughly before use

**FOR USE ON DOGS**

TO CONTROL *Fleas, Ticks* and *Lice* Starting at the tail move the dispenser rapidly and make sure that the animal's entire body is covered including the legs and underbody While spraying fluff the hair so that the spray will penetrate to the skin Make sure spray wets thoroughly but do not saturate animal Do not spray into eyes or face Do not make contact with genitalia Use protective gloves or mitts when applying product DO NOT USE ON CATS OR KITTENS

**OUTDOOR APPLICATION**

TO CONTROL *Ticks Including Ticks Which May Carry Lyme Disease* (perimeter treatment)  
Use Eagles 7 Flea & Tick Spray as a residual treatment to control *Ticks* including the *Deer Tick (Ixodes dammini)*, *Lone Star Tick (Amblyomma americanum)*, *Brown Dog Tick (Rhipicephalus sanguineus)*, *American Dog Tick (Dermacentor variabilis)* and *Western Black Legged Tick (Ixodes pacificus)* For each use area listed below apply until surface is wet (approximately 1 gal/750 1000 ft<sup>2</sup>)

Treat around windows and doors porches screens eaves patios garages under stairways and in crawl spaces where these pests may enter

Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control *Ticks* occurring there

To help control *Ticks* in outdoor activity areas treat a band of vegetation and leaf litter 6 10 feet wide and adjacent to activity areas especially where dense vegetation occurs

**FOR CONTROL OF ORNAMENTAL GARDEN INSECTS**

For use on roses dahlias asters and other ornamentals to kill *Japanese Beetles* hit by spray as well as *Whiteflies, Aphids, Exposed Thrips, Spider and Red Mites* and *Leafminers* May be used on plants as a multipurpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas

6/11

May also be used on other plants such as African violets azaleas begonias camellias carnations chrysanthemums delphiniums dogwoods English ivy Eunonymus fuschia geraniums crassula grape vines Kentia palms laurels margolds rhododendrons rubber plants snapdragons stocks wandering Jews and zinnias

Effective when used as directed against *Whiteflies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms* When spraying plants do not operate closer than 18 inches Use sweeping motion Be cautious about wetting tender foliage young plants and new growth Do not spray plants when temperatures exceed 90°F Contact insects directly whenever possible Repeat treatments as infestations occur

#### STORAGE AND DISPOSAL

**Pesticide Storage** Store in original container in a cool dry area that is inaccessible to children and pets Protect from freezing

**Pesticide Disposal** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

**Container Handling** Non refillable container Do not reuse or refill this container *If empty* Place in trash or offer for recycling if available

*If partly filled* Call your local solid waste agency for disposal instructions Never place unused product down any indoor or outdoor drain

**NOTICE** To the extent consistent with applicable law buyer assumes all risks of use storage or handling of this product not in strict accordance with directions given herewith

**Bacon Products corporation**  
P O Box22187  
Chattanooga TN 37422 2187

7/11

**SUB-LABEL B – COMMERCIAL/OCCUPATIONAL**

# Eagles-7 Flea & Tick Spray

- *[For Use Both Indoors And Outdoors*
- *A Ready-To Use Permethrin Stable Emulsion Insecticide*
- *For Control Of Fleas, Brown Dog Ticks And Lice On Premises And On Dogs*
- *For Control Of Ticks Which May Cause Lyme Disease On Premises*
- *Controls Fleas And Ticks On Dogs For 35 Days*
- *Aids in Control of Ticks on Dogs for an additional 2 weeks*
- *For control of flies and lice on livestock and premises]*

**ACTIVE INGREDIENT**

Permethrin 0.50%

**OTHER INGREDIENTS**

99.50%

100.00%

Cis/trans ratio min 35% (+/-) cis and max 65% (+/-) trans

**EPA REG NO 3487 27**

**EPA EST NO**

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See back/side panel for additional precautions

NET CONTENTS

**READ ENTIRE LABEL BEFORE EACH USE  
PET USE APPLIES TO USE ONLY ON DOGS**

**PRECAUTIONARY STATEMENTS  
Hazards To Humans & Domestic Animals**

**CAUTION**

**Personal Protective Equipment (PPE)**

Applicators and other handlers must wear  
long sleeve shirt and long pants  
socks and shoes  
chemical resistant gloves made of any water proof material (selection category A)

**User Safety Requirements**

Follow manufacturer s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product s concentrate. Do not reuse them.

**User Safety Recommendations**

User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Use Restrictions & Precautions**

- Except when applying to pets or livestock, do not enter or allow adults, children, or pets to enter the treated area until sprays have dried.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Not for broadcast use on indoor residential surfaces.
- Cover all food processing surfaces and utensils during treatment or thoroughly wash before use. Cover or remove exposed food/feed.
- Do not contaminate food or feedstuffs.



9/11

- When used in dairy barns or facilities Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects Remove or cover milking utensils before application Wash teats of animals before milking
- Remove pets and birds and cover fish aquaria before spraying and surface applications
- Do not use on electrical equipment or on surfaces that can be harmed or stained by water
- Application is prohibited directly into sewers or drains or to any area like a gutter where drainage to sewers storm drains water bodies or aquatic habitat can occur Do not allow the product to enter any drain during or after application
- For pet care Do not apply to dogs less than 12 weeks old Consult a veterinarian before applying this product on medicated debilitated aged pregnant or nursing animals Sensitivities may occur after using ANY pesticide product for pets If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water If signs continue consult a veterinarian immediately
- After applications outdoors do not water the treated area to the point of run off
- Do not make outdoor applications during rain
- Not for use in outdoor residential misting systems
- All outdoor applications must be limited to spot or crack and-crevice treatments only except for the following permitted uses
  - 1) Treatment to soil or vegetation around structures
  - 2) Applications to lawns turf and other vegetation
  - 3) Applications to building foundations up to a maximum height of 3 feet
- Other than applications to building foundations all outdoor applications to impervious surfaces such as sidewalks driveways patios porches and structural surfaces (such as windows doors and eaves) are limited to spot and crack and crevice applications only

Ready to use Turn nozzle on trigger sprayer to desired pattern Do not transfer to another container

#### SHAKE WELL BEFORE USING

#### INDOOR APPLICATION

For spot application only Do not use as a space spray indoors

#### SPOT AND CRACK & CREVICE APPLICATION

To control of *Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles* Direct the spray into hiding places cracks and crevices and behind shelves and drawers For *Silverfish*, spray bookcases For *Ants*, spray trails nests and points of entry

TO CONTROL *Carpet Beetles* Spray edges of carpeting and under carpeting and rugs Make localized application to floor and baseboards Spray directly into cracks crevices and infested areas of shelving

TO CONTROL *Bedbugs* Spray mattresses lightly particularly around tufts and seams Take beds apart and spray in all joints Treat baseboards moldings and floors

TO CONTROL *Fleas (adults and larvae)* and *Ticks, including Ticks which may carry Lyme disease (adults and larvae)* Thoroughly spray infested areas pet beds resting quarters nearby cracks and crevices along and behind baseboards moldings window and door frames and entire areas of floor and floor covering Place fresh bedding in animal quarters following treatment Allow to dry thoroughly before allowing pets to contact treated areas

TO CONTROL *Clothes Moths* Remove any infested articles from storage brush thoroughly and air for several hours in sunlight if possible Apply spray liberally to empty chests closets bureaus and other storages directing the spray into cracks joints and crevices After airing infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining

TO KILL *Lice and Louse Eggs* Spray in an inconspicuous area to test for possible staining or discoloration Inspect again after drying then proceed to spray entire area to be treated Spray from a distance of 8 to 10 inches Treat only those garments and parts of bedding including mattresses and furniture that cannot be either laundered or dry cleaned Allow all treated articles to dry thoroughly before use

10/11

**ON LIVESTOCK**

TO PROTECT CATTLE (BEEF AND DAIRY) GOATS SHEEP HOGS AND HORSES FROM *Horn Flies*, *Face Flies*, *Houseflies*, *Mosquitoes* and *Gnats* Apply a light mist sufficient to wet the surface of the hair For *Face Flies* spray face and head but do not spray into eyes

TO CONTROL *Stable Flies*, *Horse Flies* and *Deer Flies* Apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly Repeat treatment daily or at intervals necessary to give continued protection

TO CONTROL *Blood Sucking Lice* Apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair Repeat every 3 weeks if required

TO CONTROL *Poultry Lice* Spray roosts walls and nests or cages thoroughly Then spray over the birds with a fine mist

TO CONTROL *Bedbugs* and *Mites* in poultry houses Spray crevices of roost poles crack in walls and cracks in nests where the *Bedbugs* and *Mites* hide

**FOR USE ON DOGS**

TO CONTROL *Fleas*, *Ticks* and *Lice* Starting at the tail move the dispenser rapidly and make sure that the animal's entire body is covered including the legs and underbody While spraying fluff the hair so that the spray will penetrate to the skin Make sure spray wets thoroughly but do not saturate animal Do not spray into eyes or face Do not make contact with genitalia Use protective gloves or mitts when applying product **DO NOT USE ON CATS OR KITTENS**

**OUTDOOR APPLICATION**

TO CONTROL *Ticks* Including *Ticks Which May Carry Lyme Disease* (perimeter treatment)  
Use Eagles 7 Flea & Tick Spray as a residual treatment to control *Ticks* including the *Deer Tick (Ixodes dammini)*, *Lone Star Tick (Amblyomma americanum)*, *Brown Dog Tick (Rhipicephalus sanguineus)*, *American Dog Tick (Dermacentor variabilis)* and *Western Black Legged Tick (Ixodes pacificus)* For each use area listed below apply until surface is wet (approximately 1 gal/750 1000 ft<sup>2</sup>)

Treat around windows and doors porches screens eaves patios garages under stairways and in crawl spaces where these pests may enter

Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control *Ticks* occurring there

To help control *Ticks* in outdoor activity areas treat a band of vegetation and leaf litter 6 10 feet wide and adjacent to activity areas especially where dense vegetation occurs

**FOR CONTROL OF ORNAMENTAL GARDEN INSECTS**

For use on roses dahlias asters and other ornamentals to kill *Japanese Beetles* hit by spray as well as *Whiteflies*, *Aphids*, *Exposed Thrips*, *Spider* and *Red Mites* and *Leafminers* May be used on plants as a multipurpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas

May also be used on other plants such as African violets azaleas begonias camellias carnations chrysanthemums delphiniums dogwoods English ivy Eunonymus fuschia geraniums crassula grape vines Kentia palms laurels marigolds rhododendrons rubber plants snapdragons stocks wandering Jews and zinnias

Effective when used as directed against *Whiteflies*, *Aphids*, *Spider* and *Red Mites*, *Exposed Thrips*, *Lace Bugs*, *Japanese Beetles*, *Pavement Ants*, *Clover Mites* and *Armyworms* When spraying plants do not operate closer than 18 inches Use sweeping motion Be cautious about wetting tender foliage young plants and new growth Do not spray plants when temperatures exceed 90°F Contact insects directly whenever possible Repeat treatments as infestations occur

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**Pesticide Storage** Store in original container in a cool dry area that is inaccessible to children and pets Protect from freezing

11/11

**Pesticide Disposal** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

**Container Handling** Non refillable container Do not reuse or refill this container *If empty* Place in trash or offer for recycling if available

*If partly filled* Call your local solid waste agency for disposal instructions Never place unused product down any indoor or outdoor drain

NOTICE To the extent consistent with applicable law buyer assumes all risks of use storage or handling of this product not in strict accordance with directions given herewith

**Bacon Products corporation**  
P O Box22187  
Chattanooga TN 37422 2187