

ACCEPTED

APR 11 1975

3509-91

**DIRECTIONS**

**APPLICATION:** Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control. Apply dust to upper and lower surface of leaves.

**VEGETABLES**

**CORN, EGGPLANT, PEPPER, TOMATO:** To control Corn Earworm, Tomato Fruitworm, European Corn Borer, Sap Beetles, Fall Army Worms, Flea Beetles, Leafhopper, Tomato Hornworm, and Stink Bugs—apply at a rate not to exceed 20 pounds per acre. For better control of Corn Earworm apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. Reapply immediately after rain. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

**BEANS:** To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Flea Beetles, Japanese Beetles, Corn Earworm, Lygus Bug and Lima Bean Pod Borer—apply at a rate not to exceed 20 pounds per acre.

**CUCUMBER, SUMMER SQUASH:** To control Pineworm, Melonworm, Cucumber Beetles, Squash Bug and Flea Beetles—apply at a rate not to exceed 10 pounds per acre. Apply to Cucumber and Summer Squash only when plants are dry.

**POTATO:** To control Colorado Potato Beetles, Leafhoppers, Flea Beetles and Lygus Bug—apply at a rate not to exceed 10 pounds per acre.

APPLICATIONS TO ABOVE VEGETABLES CAN BE MADE UP TO DAY OF HARVEST.

**LAWNS**

To control Leafhoppers, Fall Army Worms and Sod Webworms—apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

**POULTRY**

Litter Treatment—apply 1/2 pound to 40 square feet to control Northern Fowl Mite, Chicken Mite, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand or apply by hand duster. Force dust into cracks and crevices. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

**DOGS & CATS**

To control Brown Dog Ticks and Fleas, apply liberally to animal, rub thoroughly into fur, apply to feet, between toes and on ears. Dust in and around sleeping quarters, all areas where ticks and fleas occur. Do not treat puppies or kittens less than four weeks old. Do not use on nursing puppies. Do not get powder in eyes and mouth of animals during treatment. Do not repeat within 1 week of each treatment.

**WARNING: HARMFUL IF SWALLOWED OR INHALED.** Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Remove and wash contaminated clothing with soap and hot water. Avoid contamination of feed and foodstuffs. Do not reuse container. Destroy when empty. This product is highly toxic to bees exposed to direct treatment or residues on crops. For more information may be obtained from your Cooperative Agricultural Extension Service.




10%

# SEVIN DUST

READY TO USE ON:

**VEGETABLES • GARDENS • LAWNS  
POULTRY • DOGS • CATS**



2510 E. 5th St. **SAFEGWAY FARM PRODUCTS CO., INC.** AUSTIN, TEXAS



Faint, illegible text, possibly a header or address line.

Faint, illegible text, possibly a list or description.

Faint, illegible text, possibly a list or description.

Faint, illegible text, possibly a list or description.

Faint, illegible text, possibly a list or description.

Handwritten mark or stamp, possibly a date or initials.



VEGETABLES • GARDEN  
POULTRY DOGS



25th & 5th St. SAFEWAY FARM PRODUCTS CO., INC. AUSTIN, TEXAS