

SANI-SAN

DANGER

Keep Out Of Reach Of Children. Corrosive. Causes eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures to prevent emulsification, shock, respiratory depression and convulsion may be needed. Rinse empty container thoroughly with water and discard.

FOR SALE, FOR USE BY, AND STORAGE
BY MAINTENANCE PERSONNEL ONLY

DISINFECTANT DEODORIZER SANITIZER

ACTIVE INGREDIENTS:	
n-Alkyl (60% C14, 30% C16, 5% C12, 5% C13)	
dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C12, 32% C14) dimethyl	
ethylbenzyl ammonium chlorides	5%
INERT INGREDIENTS:	90%
SANI-SAN is a concentrate and should be diluted before using.	100%

NET CONTENTS: _____ EPA Reg. No. 3525-46
EPA Est. No. 3525-NJ-1

DANGER

Keep Out Of Reach Of Children. See First Aid and other precautionary statements on left side panel.

UTICO CORPORATION
6th AVENUE & WAIT STREET
PATERSON, NEW JERSEY 07524

USE DIRECTIONS:

Household
For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For schools, homes, locker rooms, garbage pits, sink tops, corridors, classrooms, offices and shower stalls. Apply solution with mop or cloth. Sanitize with 1 ounce SANI-SAN to 4 gallons of water. Disinfect with 3 ounces SANI-SAN to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a precleaning step is recommended.

Hospitals
For floors, walls and other hard surfaces. In nursing homes, institutions, as well as sink tops, garbage pits, telephones and restrooms, use SANI-SAN at 3 1/2 ounces per 5 gallons of water. At this level SANI-SAN will be effective against Pseudomonas aeruginosa PRD-10.

Equipment
For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1/2 ounce SANI-SAN per gallon of water for 10 minutes.

RESIDUAL BACTERIOSTAT and RESIDUAL SELF-SANITIZER UNDER CONDITIONS OF HIGH RELATIVE HUMIDITY OR WET CONTAMINATION.
Fabric treated as directed will develop residual bacteriostatic activity and residual self-sanitizing activity against Gram positive (S. aureus) and Gram negative (E. coli) organisms, under conditions of high relative humidity or wet contamination.

SANI-SAN prevents the growth of ammonia producing organisms (B. ammoniagenes) which act on urine in the diapers to produce ammonia.

USE DIRECTIONS: Use 2 ounces SANI-SAN per 8 pounds of laundry in RINSE CYCLE.