

UTISEPT

Cleaner-Disinfectant-Deodorizer-Fungicide-Virucide\*  
for Hospital and Institutional Use

pg. 20/2  
PM 31  
EPA # 3525-67  
CLF 1/24/90

AOAC Phenol Coefficients

Staphylococcus aureus (ATCC No. 6538).....	104.0
Salmonella typhosa (ATCC No. 6539).....	40.6

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride.....	3.750%
Dioctyl dimethyl ammonium chloride.....	1.875%
Didecyl dimethyl ammonium chloride.....	1.875%
Alkyl(C14-50%, C12-40%, C16-10%) dimethyl benzyl ammonium chloride.....	5.000%
INERT INGREDIENTS.....	87.500%
Total	100.000%

KEEP OUT OF REACH OF CHILDREN  
DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, raw egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. Reg. No. 3525-67

E.P.A. Est. No. 3525-NJ-1

NET CONTENTS: 1 Gallon (3.785 liters)

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Add 1/2 ounce (1 tablespoon) per gallon of water.

Apply UTISEPT to previously cleaned hard surfaces, such as walls, floors, tables, chairs and bedframes with a mop or cloth.

At 1/2 ounce (1 tablespoon) per gallon use level, UTISEPT is effective against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, and Trichophyton interdigitale, the athlete's foot fungus. Germicidal performance against the three organisms has been confirmed by the AOAC Use-Dilution test. Fungicidal performance against T. interdigitale was determined by the AOAC Fungicidal test.

The broad spectrum effectiveness of UTISEPT shown by its germicidal action against the following additional organisms: Escherichia coli, Klebsiella pneumoniae, Aerobacter aerogenes (enterobacter), Salmonella schottmuelleri, Streptococcus faecalis, Shigella dysenteriae, and Brevibacterium ammoniagenes.

Not Reviewed. Registrant claims to be in accordance with  
Registration of Pesticide Products for Phase 2  
of the Federal Insecticide, Fungicide, and Rodenticide Act in Status.

\* At 1/2 ounce (1 tablespoon) per gallon use level, UTISEPT is virucidal against Herpes Simplex (a member of the virus family that causes infectious mononucleosis), Vaccinia (representative of the pox viruses), and Influenza A2 as represented by the strains commonly called the Hong Kong flu and the London flu virus, on inanimate environmental surfaces.

Cross infection is of major housekeeping concern not only in hospitals, but schools, institutions and industry. UTISEPT is formulated for this problem area. It both cleans and disinfects effectively and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection from environmental surfaces.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**

Keep Out of Reach of Children. May cause severe eye irritation or eye damage. Causes skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

**STORAGE AND DISPOSAL**

- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

**PESTICIDE DISPOSAL**

PESTICIDE, OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.


**CONTAINER DISPOSAL**

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

**GENERAL**

CONSULT FEDERAL, STATE AND LOCAL AUTHORITIES FOR APPROVED ALTERNATE PROCEDURES SUCH AS LIMITED OPEN BURNING.

UTIKEM PRODUCTS  
Div. of Qualco, Inc.  
225 Passaic Street  
Passaic, NJ 07055



**Cleaner-Disinfectant-Deodorizer-Fungicide-Virucide\***  
for Hospital and Institutional Use

PM 31  
EPA# 3525-67  
CLF 1/24/90

AOAC Phenol Coefficients

Staphylococcus aureus (ATCC No. 6538).....	104.0
Salmonella typhosa (ATCC No. 6539).....	40.6

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride.....	3.750%
Dioctyl dimethyl ammonium chloride.....	1.875%
Didecyl dimethyl ammonium chloride.....	1.875%
Alkyl(C14-50%, C12-40%, C16-10%) dimethyl benzyl ammonium chloride.....	5.000%
<u>INERT INGREDIENTS</u> .....	87.500%
Total	<u>100.000%</u>

KEEP OUT OF REACH OF CHILDREN  
DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.  
If swallowed, drink promptly a large quantity of milk, raw egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. Reg. No. 3525-67

E.P.A. Est. No. 3525-NJ-1

NET CONTENTS: 1 Gallon (3.785 liters)

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


Add 1/2 ounce (1 tablespoon) per gallon of water.

Apply UTISEPT to previously cleaned hard surfaces, such as walls, floors, tables, chairs and bedframes with a mop or cloth.

At 1/2 ounce (1 tablespoon) per gallon use level, UTISEPT is effective against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, and Trichophyton interdigitale, the athlete's foot fungus. Germicidal performance against these organisms has been confirmed by the AOAC Use-Dilution test. Fungicidal performance against T. interdigitale was determined by the AOAC Fungicidal test.

The broad spectrum effectiveness of UTISEPT shown by its germicidal action against the following additional organisms: Escherichia coli, Klebsiella pneumoniae, Aerobacter aerogenes (enterobacter), Salmonella schottmuelleri, Streptococcus faecalis, Shigella dysenteriae, and Brevibacterium ammoniagenes.

Not Reviewed. Registrant claims to be in accordance with  
Federal Register Guidance for Phase 2  
Registration, Filing to Avoid Change in Status.



\* At 1/2 ounce (1 tablespoon) per gallon use level, UTISEPT is virucidal against Herpes Simplex (a member of the virus family that causes infectious mononucleosis), Vaccinia (representative of the pox viruses), and Influenza A2 as represented by the strains commonly called the Hong Kong flu and the London flu virus, on inanimate environmental surfaces.

Cross infection is of major housekeeping concern not only in hospitals, but schools, institutions and industry. UTISEPT is formulated for this problem area. It both cleans and disinfects effectively and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection from environmental surfaces.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**

Keep Out of Reach of Children. May cause severe eye irritation or eye damage. Causes skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

**STORAGE AND DISPOSAL**

- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

**PESTICIDE DISPOSAL**

PESTICIDE, OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.


**CONTAINER DISPOSAL**

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

**GENERAL**

CONSULT FEDERAL, STATE AND LOCAL AUTHORITIES FOR APPROVED ALTERNATE PROCEDURES SUCH AS LIMITED OPEN BURNING.

UTIKEM PRODUCTS  
Div. of Qualco, Inc.  
225 Passaic Street  
Passaic, NJ 07055



Cleaner-Disinfectant-Deodorizer-Fungicide-Virucide\*  
for Hospital and Institutional Use

AOAC Phenol Coefficients

Staphylococcus aureus (ATCC No. 6538).....	104.0
Salmonella typhosa (ATCC No. 6539).....	40.6

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride.....	3.750%
Diocetyl dimethyl ammonium chloride.....	1.875%
Didecyl dimethyl ammonium chloride.....	1.875%
Alkyl(C14-50%, C12-40%, C16-10%) dimethyl benzyl ammonium chloride.....	5.000%
<u>INERT INGREDIENTS</u> .....	87.500%
Total	100.000%

KEEP OUT OF REACH OF CHILDREN  
DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, raw egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. Reg. No. 3525-67

E.P.A. Est. No. 3525-NJ-1

NET CONTENTS: 1 Gallon (3.785 liters)

DIRECTIONS FOR USE

GENERAL CLASSIFICATION


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Add 1/2 ounce (1 tablespoon) per gallon of water.

Apply UTISEPT to previously cleaned hard surfaces, such as walls, floors, tables, chairs and bedframes with a mop or cloth.

At 1/2 ounce (1 tablespoon) per gallon use level, UTISEPT is effective against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, and Trichophyton interdigitale, the athlete's foot fungus. Germicidal performance against the three organisms has been confirmed by the AOAC Use-Dilution test. Fungicidal performance against T. interdigitale was determined by the AOAC Fungicidal test.

The broad spectrum effectiveness of UTISEPT shown by its germicidal action against the following additional organisms: Escherichia coli, Klebsiella pneumoniae, Aerobacter aerogenes (enterobacter), Salmonella schottmuelleri, Streptococcus faecalis, Shigella dysenteriae, and Brevibacterium ammoniagenes.



Not Reviewed. Registrant claims to be in accordance with  
Federal Regulation of Pesticide Product Guidance for Phase 2  
of the Act, Part 2.12, Active to Inert Change in Status.

\* At 1/2 ounce (1 tablespoon) per gallon use level, UTISEPT is virucidal against Herpes Simplex (a member of the virus family that causes infectious mononucleosis), Vaccinia (representative of the pox viruses), and Influenza A2 as represented by the strains commonly called the Hong Kong flu and the London flu virus, on inanimate environmental surfaces.

Cross infection is of major housekeeping concern not only in hospitals, but schools, institutions and industry. UTISEPT is formulated for this problem area. It both cleans and disinfects effectively and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection from environmental surfaces.

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Keep Out of Reach of Children. May cause severe eye irritation or eye damage. Causes skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL

PESTICIDE, OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE AND LOCAL AUTHORITIES FOR APPROVED ALTERNATE PROCEDURES SUCH AS LIMITED OPEN BURNING.

UTIKEM PRODUCTS  
Div. of Qualco, Inc.  
225 Passaic Street  
Passaic, NJ 07055

