

HIDE

SEVEN DUST 5%

ACTIVE INGREDIENT: Carbaryl (1 Naphthyl N Methylcarbamate)	5.0%
INERT INGREDIENTS:	95.0%
TOTAL	100.0%

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid contamination of food or foodstuffs.

NOTE TO PHYSICIANS: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Dust the upper and lower surfaces of the foliage thoroughly.

BEANS. Against Mexican bean beetle, leafhoppers, lygus, and stink bugs, dust thoroughly, beginning when insects first appear and repeat every 7 to 10 days or as needed. Apply 1 to 2 oz. per 50 foot row.

CUCUMBER, SUMMER SQUASH: Against striped and spotted cucumber beetles, squash bugs, flea beetles, pickleworms use 1 to 2 oz. per 50 foot row. To avoid possible injury to tender foliage do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

TOMATO, EGGPLANT, PEPPER: Against tomato fruitworm, fall armyworm, and flea beetle, dust thoroughly when insects first appear and every 7 to 10 days if necessary. Use 1 to 2 oz. per 50 foot row.

POTATO: Against Colorado potato beetle, leafhopper and flea beetle apply 1 to 2 oz. per 50 foot row.

COLLARDS, TURNIPS (tops), MUSTARD GREENS. Against flea beetle, corn earworm, armyworms, imported cabbage worm use 1 to 2 oz. per 50 foot row when insects first appear and repeat every 7 to 10 days if necessary for control.

DOGS AND CATS: For Fleas and Ticks apply dust liberally over entire animal, avoiding eyes, and rub thoroughly in to skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if necessary. Avoid treatment of kittens less than 4 weeks of age.

Dust liberally in and around sleeping quarters, kennels, and dog dust into cracks and crevices. Repeat as necessary. Do not use this product on poultry. Do not reuse container.

DISPOSAL: Do not reuse empty container. Wrap in newspaper and put in trash collection.

EPA EST NO 3325-1A-1
EPA REG. NO. 3325-72

Net Wt. 16 oz.

HIDE INTERNATIONAL
Box 449
Leon, Iowa 50144

