

3536-4

07/08/1963

111

DIRECTIONS

RATS

Suggested use: Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

MICE

Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

NEW IMPROVED

H-K RAT BAIT

For Hard to Kill Rats & Mice

A better and more effective rodenticide.

The most notable advance in rodenticides since Warfarin.

The antibacterial agent increases the effectiveness of the product and assures thorough control of rats and mice. Especially recommended for rat and mouse infestations which are difficult to control.




a better, more effective warfarin rodenticide

ACTIVE INGREDIENTS:

Sulfaquinoxaline (N1-(2-quinoxaliny) sulfanilamide)	0.025%
Warfarin 3(a-acetonylbenzyl)-4-hydroycoumarin	0.025%

INERT INGREDIENTS: 99.950%



* PROLIN is a Trademark of the Wisconsin Alumni Research Foundation

GUARANTEE — If fully satisfactory results are not obtained, your money will be refunded upon return of the unused portion to

PIPESTONE PRODUCTS

PIPESTONE, MINNESOTA

NET WEIGHT: 5 LBS.

ACCEPTED
JUL 8, 1963
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3536-4