



ACCEPTED

FEB 18 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 3536-4



KILLS NORWAY RATS,  
ROOF RATS  
AND HOUSE MICE

## ACTIVE INGREDIENTS:

Warfarin, 3 - (a-acetylbenzyl) - 4-hydroxycoumarin .....	0.025%
INERT INGREDIENTS .....	99.975%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN

## CAUTION

PRECAUTIONARY STATEMENTS:  
HAZARD TO HUMAN & DOMESTIC ANIMALS

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

**NOTE TO PHYSICIAN -** If ingested, administer Vitamin K, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

## ENVIRONMENTAL HAZARDS

KEEP OUT OF LAKES, STREAMS OR PONDS

**STORAGE:** Store only in original container, in a dry place inaccessible to children and pets.

**DISPOSAL:** Do not reuse empty container. Securely wrap in newspaper and discard in trash.

## DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT  
IN A MANNER INCONSISTENT WITH ITS LABELING

**USE RESTRICTIONS -** For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS -** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

## APPLICATION DIRECTIONS

**RATS:** Place 4 to 16 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped.

**MICE:** Apply ¼ to ½ ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

E.P.A. REG. NO. 3536-4

E.P.A. Est. No. 3536-MN-02

**PIPESTONE PRODUCTS CO., INC.**  
TROSKY, MINNESOTA 56177

NET CONTENTS 5 LBS.