

**LYNWOOD LABORATORIES, INC.**

945 GREAT PLAIN AVENUE • NEEDHAM, MA • 02492
TEL: (781) 449-6776 • FAX: (781) 444 2454 • EMAIL: irvinglk@aol.com

March 1, 2006

NOTIFICATION**MAR 29 2006**

Ms. Sherada Hobgood
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
US Environmental Protection Agency
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202-4501

Dear Ms. Hobgood:

In accordance with EPA procedure, we are pleased to submit notification of our Shoo-fly Roach Frase II

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

Sincerely,

Paul G. Ledoux
National Sales Manager
Lynwood Laboratories
Tel: 781-449-6776

PGL:cg

Cc: I. Lynwood Kanin, President, Lynwood Laboratories
Cc: Warren Bennett, Shield Packaging

SHOOFLY®

The Best Products For Insect Control

APIA

DIRECTIONS: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling: TO OPERATE: For crack and crevice injection use. Remove protective cap. Hold container upright. Attach injection tube and insert into cracks and crevices and tunnels where insects may be hiding or breeding. Place injection tip 1/2 inch into cracks, crevices, holes and other small openings where insects may be a problem. Release a one second spray per spot. In serious infestations, if possible, place spots approximately 12 inches apart. In light infestations or for preventative treatment, place spots approximately three feet apart. In long running cracks, move the injector tip along the crack while spraying. Move at a rate of one linear foot per second in heavy infestations and three feet per second in light infestations. See the additional directions below for effective control of specified insects.

INDOOR USE: LOCALIZED TREATMENT OF WOOD INFESTING TERMITES, CARPENTER ANTS, CARPENTER BEES, WOOD BORERS, WOOD WASPS: Locate area of infestation and/or insect tunnels and cavities. Using a crack & crevice tip, spray into tunnels, galleries and inaccessible closed voids. In infested areas, small holes should be drilled eight to ten inches apart to insure thorough insecticide gallery treatment. Be sure not to contact electrical wiring or plumbing when treating. For Carpenter Bees, seal the entrance holes with small pieces of stainless steel wool after application. The Shoo-fly spray protects up to four weeks.

The purpose of such applications of Shoo-Fly Termite and Carpenter Ant killer for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. For active termite infestations, get a professional inspection.

Ants: Spray freely around ant trails, doors and windows and other places where ants enter premises around nests hidden under steps, brickwork, concrete, etc. Break apart accessible nests and spray freely on and around debris. Repeat every four weeks. **Carpenter Ants:** For effective control, locate and treat nests and surrounding areas. Apply around doors and windows and other places where ants enter premises and where they crawl and hide. Inject into wall voids and infested wood through existing openings or newly drilled small holes. **Outdoors:** Pests on Outside Surfaces Of Buildings: For control of building infestation due to Ants, Clover Mites, Crickets, Earwigs, Millipedes and Sowbugs (Pillbugs). Spray foundation of building where insects are active and may find entrance up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include (but are not limited to) screens, window frames, eaves, porches, patios, garage and refuse dumps. Repeat treatment every 4 weeks.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS:
CAUTION: Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

SHOO-FLY Blast Termite and Carpenter Ant Killer may be used as a crack and crevice treatment or as a spot application in non-food/feed areas. All areas where insects hide or through which insects may enter should be treated. Not for use in Federally Inspected Meat and Poultry Plants. Do not apply this product in occupied patient rooms or in any rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Storage: Do not store near heat or open flame. Store in cool dry area away from children. Container Disposal: Replace cap, wrap container in several layers of newspaper and discard in trash. Do not incinerate or puncture. Can may be disposed of with other recyclables where local law permits.

EPA Est. No. 5693-MA-1
EPA Reg. No. 3546-37

Manufactured for LYNNWOOD LABORATORIES
Needham, Massachusetts 02492
Fax: (781)-444-2454

Contains no CFC's or other ozone depleting substances, federal regulations prohibit CFC propellants in aerosols




ACCEPTED
JAN - 5 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for use pesticides registered under EPA Reg. No. 3546-37

NOTIFICATION


MAR 2 9 2006

REV. 10/05



shoo-fly

Termite & Carpenter Ant Killer



KILLS Termites, Carpenter Ants and Other Wood Infesting Insects

KEEP OUT OF REACH OF CHILDREN
CAUTION
CONTENTS UNDER PRESSURE. SEE ADDITIONAL CAUTIONS ON BACK PANEL
NET CONTENTS 15 OZ.

228

IDENTIFICATION

MAR 29 2006

FRONT OF ROACH ERASE II LABEL

SHOO-FLY

BLAST

ROACH ERASE II

Instant Kill
Plus Residual
Kills

Ants, Cockroaches,
Fleas, Termites, Ticks,
Spiders, Crickets, Scorpions,
Carpenter Bees

ACTIVE INGREDIENTS:

• Tetramethrin.....	0.150%
** Permethrin.....	0.375%
*** Piperonyl Butoxide Technical.....	0.750%
INERT INGREDIENTS	<u>98.725%</u>
	100.00%

- 1-cyclohexene 1,2 dicarboximidomethyl-3-(2methlpropenyl) cyclopropanecarboxlate
- ******(3-phenoxyphenyl) methyl (+/-) cis/trans 3-)2,2 dichloroethenyl) 2,2-dimethylcylopropanecarboxlate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.
- ******* Equivalent to min. 0.60% (butylcarbityl)(6-propylpiperonyl) ether and 0.15% related compounds
- TETRAPEM - Registered Trademark of AgreEvo Environmental Health., Inc.

4 7 8

KEEP OUT OF REACH OF CHILDREN

CAUTION

CONTENTS UNDER PRESSURE
SEE ADDITIONAL CAUTIONS ON BACK PANEL

NET CONTENTS 10 oz.

1 1
1 1 1 1

Roach Erase II label left side

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using.

Do not allow children or pets into the treated area until dry.

REDISUAL ACTION

One spraying of Shoo-Fly Roach Erase II lasts 4 weeks. When used as directed this spray leaves an invisible film that continues to kill roaches, ants, carwigs, spiders, and flies.

INDOOR USE

ROACHES, SILVERFISH, CRICKETS: Spray heavily into all cracks, around baseboards and other places where these insects hide. Also spray until wet, surfaces behind and beneath cabinets, refrigerators, sinks, stoves, etc. Spray other localized areas infested with pests. Repeat treatment as required.

SPIDERS: Spray spiders and their immediate vicinity thoroughly. Do not spray inside cabinets where food is stored. Discard infested food packages. Do not spray across top of stove or near pilot lights.

ANTS: Spray freely around ant trails, doors and windows and other places where ants enter premises around nests hidden under steps, brickwork, concrete, etc. Break apart accessible nests and spray freely on and around debris. Repeat every four weeks.

CARPENTER ANTS: For effective control, locate and treat nests and surrounding areas. Apply around doors and windows and other places where ants enter premises and where they crawl and hide. Inject into wall voids and infested wood through existing openings or newly drilled small holes.

Take care not to introduce this product into the air during application. Use only in well ventilated areas. Do not apply where children are likely to frequently contact treated surfaces. Do not treat pets or plants with this product. Remove pets and cover fish bowls before spraying. Do not apply when other persons are in immediate area. Do not apply in areas, surfaces, or utensils which will contact food or feed. Exposed food should be covered or removed during application. All food processing surfaces and utensils should be covered during application or thoroughly washed with soap and water before use. Do not use in edible product areas of food processing plants, restaurants, or other places where food is commercially prepared or processed.

Do not spray floors, wallcoverings, ceilings without first testing for staining. Be sure not to contact electrical wiring or plumbing when treating.

EPA Est. No. 5693-MA-1

EPA Reg. No. 3546-37

CONTAINS NO CFC'S OR
OTHER OZONE DEPLETING
SUBSTANCES, FEDERAL
REGULATIONS PROHIBIT CFC
PROPELLANTS IN AEROSOLS

Roach Erase II label right side

OUTDOOR USE

Carpenter Ants: For effective control, locate and treat nests and surrounding areas. Apply around doors and windows and other places where ants enter premises and where they crawl and hide. Inject into wall voids and infested wood through existing openings or newly drilled small holes.

Pests on Outside Surfaces of Buildings: For control of building infestation due to Ants, Crickets, Earwigs, Millipedes and Sowbugs. Spray foundation of building where insects are active and may find entrance up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include (but are not limited to) screens, window frames, eaves, porches, patios, garage and refuse dumps. Repeat treatment every 4 weeks. Be sure not to contact electrical wiring or plumbing when treating. For Carpenter Bees, seal the entrance holes with small pieces of stainless steel wool after application.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Caution: Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses if present after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of waste.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

Do not use in food/feed areas handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be removed.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.

SHOO-FLY Roach Ease II may be used as a crack and crevice treatment or as a spot application in non-food/feed areas. All areas where insects hide or through which insects may enter should be treated. Do not apply this product in occupied patient rooms or in any rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Storage: Do not store near heat or open flame. Store in cool dry area away from children. Container Disposal: Replace cap, wrap container in several layers of newspaper and discard in trash. Do not incinerate or puncture. Can may be disposed of with other recyclables where local law permits.

WARRANTY & DISCLAIMER

Lynwood Laboratories, Inc. guarantees this product to the extent of the purchase price only when used as directed.

Manufactured for
Lynwood Laboratories, Inc.
Needham, MA 02492
781-449-6776
Fax: 781-444-2454

0 49711 991108