

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under EPA Reg. No. 3618-29
 AUG 9 1975
ACCEPTED

INDUSTRIAL Formula

20% LINDANE

DIRECTIONS: Tobacco — For the control of wireworms, use a solution containing 2-4 oz. per 100 gallons of water. Apply at the rate of 1/2 pt. per plant or 200-250 gallons of solution per acre—depending on type of soil. Do not use when rotation with root crops and peanuts.

Livestock — Beef Cattle and Hogs — Spray entire animal with solution containing 1 pt. 20% Lindane per 100 gallons of water. For dipping to control lice and ticks, submerge animal completely, including head, at least twice in solution containing 1 1/2 pts. 20% Lindane per 100 gallons water. Do not allow animals to drink dipping solution.

Do not dip animals within 60 days or spray within 30 days of slaughter. Do not treat animals under 3 months of age. Avoid treatment in cold stormy weather. Do not treat overheated or sick animals.

This product is toxic to fish, birds, and other wildlife. Keep out of streams, ponds, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas

For use in transplanting water for protection against wireworms in tobacco plants, and control of cattle horn flies, ticks and flies.

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride from Lindane 20.00%

Xylene 77.00%

INERT INGREDIENTS 3.00%

TOTAL 100.00%

CAUTION: Keep Out of Reach of Children

May cause skin and eye irritation. Can be absorbed through skin. Avoid contact with skin. Harmful if swallowed. Avoid inhaling spray mist. Avoid contamination of feed and food-stuffs. Inflammable, keep away from open flame. Do not smoke while spraying. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Antidote:

External — In case of spillage on skin, wash with soap and water. For eyes, flush with plenty of water, get medical attention.

Internal — Drink emetic of mustard and call a physician.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

manufactured by

INDUSTRIAL COLLOIDS & CHEMICALS, INC.

127 West Jackson Avenue — Knoxville, Tennessee

NET CONTENTS: 1 GAL.

treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Farm Buildings and Barns — Spray thoroughly inside and out with solution containing 1 1/2 gallons 20% Lindane per 100 gallons water. Apply at the rate of one gallon per 500 sq. ft., wetting all surfaces to the point of run-off. Protect all feed and feed troughs before spraying. Remove all animals from building. Do not contaminate milk, milking equipment, feed or drinking water. **NOTE:** Do not use treated manure on land where root crops or peanuts are to be planted.

Forage Crops — (Alfalfa and Clover)

— For control of sprouting apply 20% Lindane at the rate of 1 pt. per acre with sufficient water for treatment. Apply when forage is below 6 inches tall. Do not graze or cut alfalfa or clover for forage within 30 days after treatment.

NOTE: If you do not have a material supply to use in your area, contact your distributor.

EPA No. 3618-29

ACCEPTED WITH COMMENTS