

APPLICATION TO GRAIN SURFACES: APPLY the spray as a coarse wet spray to the surface of grain at the rate of 1 gallon to 750 square feet of top surface. At the same time spray the surface of the bin above the grain. If the grain is already infested with Indian Meal Moth, rake the surface after spraying and spray the surface again. The raking should be to a

REMOVAL OF GRAIN FROM BIN: After applying this spray to the surface of grain the top surface of the grain must be thoroughly mixed with the rest of the bin contents before loading out. Spraying of the top surface is not recommended for box cars or small grain masses. To use this spray on box cars of grain or small grain storages may result in using more than the legal tolerance of 2 ppm. of the toxicant.

WARNING: Keep out of the reach of children. Harmful if swallowed. Avoid prolonged or repeated breathing of vapor. Avoid prolonged or repeated

DO NOT FREEZE. Store in warm

NPI

MARNO

GRAIN & ELEVATOR SPRAY

ACTIVE INGREDIENTS: 100%

2.500% Technical Methoxychlor* Technical Piperonyl Butoxide** .063% .025% **Pyrethrins**

15.000% Aromatic Petroleum Derivative Solvent 82.412% Petroleum Distillate

*Equivalent to 2.2% of 2,2-bis(p-methoxyphenyl)-1,1,1trichloroethane and 0.3% of other isomers and related compounds.

**Equivalent to 0.05% of (butyt_carbityl) (6-propyl piperonyl) ether and 0.013% of related compounds.

Keep Out of Reach of Children CAUTION: en See Side Panel for Additional Cautions

Fed. Reg. No. 3624-10/

CONTENTS

GALLONS

Manufactured for

BERNARD E. FRIEL COMPANY

P.O. Box 5207, Plaza Station Kansas City 12, Missouri

H

DIC

 $\mathbf{O} \subseteq$