


3624-127



# MULTIPURPOSE CONTACT SPRAY

INDUSTRIAL SPRAY FOR FOOD PROCESSING INDUSTRIES  
FOR USE IN FOGGING EQUIPMENT OR TANK TYPE SPRAYERS

ACTIVE INGREDIENTS: 100%

Pyrethrins	0.34%
Technical Piperonyl Butoxide**	1.70%
Petroleum Distillate**	97.96%

\*EPA registered active ingredient for use in commercial agricultural spraying. Active ingredient concentration: 0.34% Pyrethrins and 1.70% Piperonyl Butoxide. EPA Reg. No. 141-127.

**CAUTION:**  
KEEP OUT OF THE REACH OF CHILDREN.

EPA Reg. No. 141-127

CONTENTS                                  GALLONS

**NOVA PRODUCTS, INC.**

Kansas City, Kansas 66119

**OBJECTIONS**

...of the spray, it is not suitable for the use of...  
...of the spray, it is not suitable for the use of...  
...of the spray, it is not suitable for the use of...

**Sanitation.** Good sanitation is necessary for insect control. Before spraying or fogging an area of a processing plant, all food ingredients, food waste, and other materials should be removed and destroyed. All food products and raw materials should be covered. Product zones should be covered. If possible, areas to be sprayed should be kept dry. A minimum of 24 hours should elapse before...

**TESTING INSECTS.** Insecticides should be tested on a small scale before being used on a large scale. The type of insecticide to be used should be determined by the type of insect. The type of insecticide to be used should be determined by the type of insect. The type of insecticide to be used should be determined by the type of insect.

...of the spray, it is not suitable for the use of...  
...of the spray, it is not suitable for the use of...

**WASHING INSECTS.** Insecticides should be tested on a small scale before being used on a large scale. The type of insecticide to be used should be determined by the type of insect. The type of insecticide to be used should be determined by the type of insect.

**CAUTION**

Insecticides should be tested on a small scale before being used on a large scale. The type of insecticide to be used should be determined by the type of insect. The type of insecticide to be used should be determined by the type of insect.

Insecticides should be tested on a small scale before being used on a large scale. The type of insecticide to be used should be determined by the type of insect. The type of insecticide to be used should be determined by the type of insect.

