

BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Do not contaminate food or feedstuffs. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not permit children or pets to go onto the sprayed grass until spray has completely dried.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to water (lakes, streams or ponds). Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal.
Pesticide Disposal: Pesticide spray mixture, or residue that cannot be used or chemically reprocessed must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.
Container Disposal: Metal (all sizes)—Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in a sanitary landfill, or by other approved State and local procedures.
 Glass (all)—Triple rinse (or equivalent) and offer for recycling, reconditioning, or by other approved State and local procedures.
Labels: Contact Federal, State, or local officials for disposal procedures for unlabelled products.

EPAA 70

01101

E-2 with HAN

| | |
|--------------------------------|--------|
| ACTIVE INGREDIENTS: | |
| *Methoxychlor, Technical | 25.0% |
| Heavy Aromatic Naphtha | 71.0% |
| INERT INGREDIENTS | |
| | 100.0% |

Equivalent to Saps 2, 2, 4, 4'-tetrahydroxy-1, 1, 1-trichloroethane
 and 3, 3, 5-trichloro-4-hydroxybenzoic acid and related compounds.

3624-171

the Federal Government has
 approved this product for use
 on the following crops:

CAUTION: IF SWALLOWED— Call a physician immediately. Get first aid information from the label. Do not induce vomiting unless a physician advises you to do so. Vomiting may cause aspiration pneumonia or other complications. If a physician is not available, give 1-2 glasses of water or milk. If a physician is available, give 1-2 glasses of water or milk. If a physician is not available, give 1-2 glasses of water or milk. If a physician is not available, give 1-2 glasses of water or milk.

IF SWALLOWED— Call a physician immediately. Get first aid information from the label. Do not induce vomiting unless a physician advises you to do so. Vomiting may cause aspiration pneumonia or other complications. If a physician is not available, give 1-2 glasses of water or milk. If a physician is available, give 1-2 glasses of water or milk. If a physician is not available, give 1-2 glasses of water or milk.

EPAN

BEST DOCUMENT AVAILABLE

E-2 with HAN

Grain and Animal Dip — 20% HAN

FIVE INGREDIENTS:

| | |
|--------------------------------|---------------|
| *Methoxychlor, Technical | 25.0% |
| Heavy Aromatic Naphtha | 71.0% |
| OT INGREDIENTS | 4.0% |
| | 100.0% |

2,2-bis (p-methoxyphenyl)-1,1,1-trifluoroethane and other isomers and related compounds.

3624-171

WARNING

Followed—Call a physician immediately if you experience any of the following symptoms: dizziness, headache, nausea, vomiting, or difficulty breathing. If you experience any of these symptoms, stop using the product immediately and seek medical attention. Do not use this product in enclosed spaces or near food or feed. Avoid contact with skin and eyes. Wash hands thoroughly after use. Do not use this product on children or animals.

Apply thoroughly and apply as dip. For use on effective for control of sheep and goat lice, apply the above solution as a dip, making sure each animal is held in vat at least 2 minutes and submerged twice during dipping.

Do not apply to lactating goats. Do not apply to dairy calves. Do not dip calves under 1 Month of age.

For use in cable-type backrubbers or automatic cattle oilers to control hornflies, and aid in the reduction of lice infestations on beef cattle, dilute 1 gallon of Methoxychlor E-2 with 4 gallons of fuel oil (makes a 5% solution.) Apply this solution to saturate the covering on the cable-type backrubbers or use to fill the reservoir on automatic type cattle oilers. Repeat every two weeks or as necessary for continued control. An occasional saturation of the covering on cable-type backrubbers with straight fuel oil after several treatments with the Methoxychlor E-2 solution may be desirable. Do not apply this solution to animals by spraying, dipping, or washing.

RESIDUAL SPRAY: In dairy barns, milk houses and animal shelters, for control of stable flies, house flies and fleas, dilute 1 gallon in 4 gallons of water and apply at the rate of 1 gallon per 500 square feet. Dairy animals should be removed from building during spraying operation and feed and water should be protected to prevent contamination. For the control of fleas apply sprays to surfaces they frequent, such as cracks in walls and floors as well as to bedding in animal's quarters. For roaches treat baseboards, under and around equipment and around plumbing fixtures. Re-treat as necessary for continued control.

Control in poultry houses: Dilute 1 gallon of Methoxychlor E-2 with 4 gallons of water and apply at the rate of 1 gallon per 500 square feet. Apply to walls, floors, and ceilings. Repeat every two weeks or as necessary for continued control.

Apply thoroughly and apply as dip. For use on effective for control of sheep and goat lice, apply the above solution as a dip, making sure each animal is held in vat at least 2 minutes and submerged twice during dipping.

Do not apply to lactating goats. Do not apply to dairy calves. Do not dip calves under 1 Month of age.

For use in cable-type backrubbers or automatic cattle oilers to control hornflies, and aid in the reduction of lice infestations on beef cattle, dilute 1 gallon of Methoxychlor E-2 with 4 gallons of fuel oil (makes a 5% solution.) Apply this solution to saturate the covering on the cable-type backrubbers or use to fill the reservoir on automatic type cattle oilers. Repeat every two weeks or as necessary for continued control. An occasional saturation of the covering on cable-type backrubbers with straight fuel oil after several treatments with the Methoxychlor E-2 solution may be desirable. Do not apply this solution to animals by spraying, dipping, or washing.

RESIDUAL SPRAY: In dairy barns, milk houses and animal shelters, for control of stable flies, house flies and fleas, dilute 1 gallon in 4 gallons of water and apply at the rate of 1 gallon per 500 square feet. Dairy animals should be removed from building during spraying operation and feed and water should be protected to prevent contamination. For the control of fleas apply sprays to surfaces they frequent, such as cracks in walls and floors as well as to bedding in animal's quarters. For roaches treat baseboards, under and around equipment and around plumbing fixtures. Re-treat as necessary for continued control.

Control in poultry houses: Dilute 1 gallon of Methoxychlor E-2 with 4 gallons of water and apply at the rate of 1 gallon per 500 square feet. Apply to walls, floors, and ceilings. Repeat every two weeks or as necessary for continued control.

NOT APPLY WITHIN SEVEN DAYS OF HARVESTING OR GRAZING: For the control of velvet bean caterpillar on soybeans, velvet beans and peanuts, apply 2 quarts per acre in sufficient water to give adequate coverage. Make application when first signs of infestation are noted. Repeat application in 10 days if needed. If two or more applications are made, allow 14 days between last application and harvest.

FOR CONTROL OF ELM BARK BEETLES (CARRIERS OF DUTCH ELM DISEASE)

DORMANT TREATMENT: Apply Methoxychlor spray (50 gallons Methoxychlor E-2 with 100 gallons of water) when using a mist blower or apply Methoxychlor spray (8 gallons Methoxychlor E-2 with 100 gallons of water) when using a hydraulic or conventional sprayer any time after the leaves drop in the fall, up to the appearance of elm flowers or leaves, usually March in the South, or April in the North. Thoroughly spray all bark surfaces. The average 50 ft. c'm will require 2 to 3 gallons of spray mixture when using a mist blower, and about 25 gallons when using a hydraulic sprayer. Spray only when temperatures are above freezing and good spraying conditions prevail.

SUMMER TREATMENT WHEN REQUIRED: Apply 600 Methoxychlor spray (25 gallons Methoxychlor E-2 with 75 gallons water) when using a mist blower or apply a 1% Methoxychlor spray (4 gallons Methoxychlor E-2 with 50 gallons water) when using a hydraulic or conventional sprayer. Repeat every two weeks or as necessary for continued control.