

13-3-71
3685-125

OXFORD IX-91 IODOPHOR CONCENTRATE

Disinfectant - Sanitizer - Detergent

Non-Staining in Use Solution

IMPORTANT—Use fresh solutions after each clean-up. The daily use of IX-91 prevents milkstone build-up. A yellow color on equipment indicates milkstone build-up and can be removed by brushing with IX 91, allowing a few minutes contact then rinsing.

GENERAL INFORMATION—IX-91 solutions wet contaminated surfaces. IX-91 solutions are self-indicating of germicidal strength. A yellow color indicates solution is germicidal. Replace solution when yellow color disappears. Clean all surfaces with suitable detergent and rinse prior to use of IX-91 solutions. Rinse or scrape soil from surfaces prior to detergent-sanitizer applications. Use a minimum exposure of two minutes at a temperature from 75° to 100° F. Use longer exposures on porous surfaces. Do not use water over 100° F.

DIRECTIONS

1. General Disinfection:
Equipment in Poultry Plant, Hatcheries, Meat Plants and Food Processing Plants. Clean with a suitable detergent and rinse thoroughly. To disinfect, use solution D for non-porous surfaces and solution E for porous surfaces. Allow a minimum of 2 minutes exposure and rinse thoroughly. Follow recommendations of Federal, State or local regulatory groups.
2. Cafeterias, Restaurants, Bars. Eating Utensils, cutlery, dishes, glassware
 - a. Scrape and pre-wash utensils and glasses.
 - b. Wash with a good detergent or compatible cleaner.
 - c. Rinse with clean water.

K-71

ACTIVE INGREDIENTS:

Nonylphenoxypoly (ethyleneoxy) ethanoliiodine complex, 16.75%; Phosphoric Acid, 6.00%; Isopropyl alcohol, 5.00%; Nonylphenoxypoly (ethyleneoxy) ethanol, 4.00%; Potassium iodide, 2.00%.

INERT INGREDIENTS: 66.25%

*Provides a minimum of 1.75% titratable iodine.

DILUTION TABLE

Solution	Dilution	Oz. per 5 gal. water	PPM Titratable Iodine
A	1:1280	1	12.5
B	1:640	1	25.0
C	1:320	2	50.0
D	1:213	3	75.0
E	1:128	5	125.0

EPA Reg. No. 3635-125

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid getting in eyes or on skin. If IX 91 gets in eyes, flush with plenty of water. If irritation persists, get medical attention.

Do not contaminate food with IX 91 or its solution. Rinse surface treated with IX-91 prior to contact with food. Avoid contact with alkalis, soap and detergents.

Do not reuse empty drum. Rinse thoroughly and discard.

Net Contents _____ Gallons

- d. Sanitize in a solution of 1 ounce to 5 gallons of water (12.5 ppm - Solution A). Immerse all utensils for at least 2 minutes or for contact time specified by governing sanitary code.
- e. Place sanitized utensils on a rack or drainboard to air dry.

3. Hospitals

- a. Surgical Instruments and Equipment. Clean and rinse articles in a suitable detergent solution. Immerse in solution E for at least 2 minutes. IX-91 will kill *Staphylococcus aureus* and many other bacteria.

4. Dairy Farm, Milking Machines and Utensils

- a. Flush all residual milk from equipment with cold or lukewarm water after each milking. Do not use hot water. Take equipment apart and immerse all parts except pulsator in solution B and thoroughly brush clean. A fresh solution B should be used to wash pails and strainers. Rinse with warm water, reassemble and rack in a clean, dust-free place.

- b. Disinfect with solution A for 2 minutes.

5. Dairy Plant: After processing milk, flush residual milk from equipment and all surfaces with lukewarm water. Wash with a suitable detergent. Rinse with water. Sanitize surfaces with solution B. Allow at least a 2 minute exposure and rinse with water. Follow recommendations of local Health Departments.

'S 6