

# OXFORD VAPOGENE

## DISINFECTANT — DETERGENT

### A.O.A.C. USE DILUTION METHOD, 11th EDITION

Staphylococcus aureus	1:256
Salmonella choleraesuis	1:256
Pseudomonas aeruginosa	1:256

VAPOGENE is a viricidal agent: against the enveloped, lipophilic viruses represented by herpesvirus (strain HF) and influenza virus A2/Aichi/2/68 (Hong Kong). VAPOGENE at a 1:256 dilution is viricidal.

VAPOGENE is Salmonellacidal — effective against the organisms which cause pullorum disease, fowl typhoid, and paratyphoid when tested using the A.O.A.C. USE-DILUTION METHOD, 11th EDITION.

Salmonella pullorum	1:256
Salmonella typhimurium	1:256
Salmonella typhosa	1:256
Salmonella choleraesuis	1:256

### DIRECTIONS

1. For disinfecting, cleaning, and deodorizing in food processing plants and equipment, use a solution of ½ ounce per gallon (1:256). Mop, wipe or swab this solution on surfaces. Leave in contact 10 minutes. Rinse all surfaces coming in contact with food with potable water before re-use.

2. For disinfecting and cleaning garbage and trash receptacles, sick room equipment, floors, walls, patient rooms, use ½ ounce of VAPOGENE to 1 gallon of water (1:256 dilution). For porous surfaces, increase concentration to 1 ounce per gallon of water.

3. For non-food disinfection in animal and poultry houses, hatcheries, stockyards, zoos, veterinary animal hospitals: Do not use in milking stalls, milk parlors or milk houses. Remove all the animals and feeds from the premises. Remove all litter and manure from floors and other surfaces, empty all water troughs and feed racks. Saturate all surfaces with a solution of ½ oz. per gallon (1:256). Ventilate buildings and other closed spaces. Do not house poultry, livestock or use equipment until treatment has been absorbed or dried. All treated feeders, fountains and waterers must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.

### ACTIVE INGREDIENTS:

Isopropanol, 12.00%; Potassium laurate, myristate and related compounds, 11.80%; Ortho-phenylphenol, 6.37%; paratertiarybutylphenol, 4.93%; 2,4,5-trichlorophenol, 2.37%; Ethylenediaminetetraacetic acid tetrasodium salt, 2.00%; 4 and 6 chloro-2 phenylphenol 1.80%; Sodium mono- and dimethyl naphthalene sulfonate, 0.50%; Essential Oils, 0.30%.

INERT INGREDIENTS ..... 57.93%

## WARNING

### KEEP OUT OF REACH OF CHILDREN

Causes severe eye or skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

### FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of water. Avoid alcohol, call a physician immediately.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

EPA Reg. No. 3635-131

A72

PHD-15

Net Content \_\_\_\_\_

4. VAPOGENE is a prophylactic agent in preventing transmission of athlete's foot fungi. Prepare a solution of 5/8 ounce per gal. of water (1:200 dilution); mop or swab this solution onto floors and other surfaces normally contacted by the feet in locker rooms, shower stalls, bathrooms, and around swimming pools.

5. VAPOGENE is conductive—may be used on conductive floorings as it does not interfere with electrical conductivity, as specified in the procedures in accordance with Paragraph 252, "Conductive Flooring of the National Board of Fire Underwriters," Specifications No. 56.

VAPOGENE is recommended for use in fogging operations to reduce airborne bacteria. Fogging with 1 ounce of VAPOGENE to 1 gallon of water 6 minutes per 1000 cubic feet of space will effectively reduce bacteria. Allow a minimum of two hours after fogging has stopped before room can be occupied.

STAPHYLOCIDAL — VIRICIDAL — FUNGICIDAL —  
PSEUDOMONACIDAL — TUBERCULOCIDAL —  
SALMONELLACIDAL — CONDUCTIVE

VAPOGENE disinfectant is a broad spectrum, powerful germicide and deodorant formulated with multiple germicidal agents to provide potent activity and detergency.

VAPOGENE is also effective against the following organisms using the A.O.A.C. USE-DILUTION METHOD, 11th EDITION

Proteus vulgaris	1:256
Streptococcus mitis	1:256
Escherichia coli	1:256

### A.O.A.C. TUBERCULOCIDAL TEST

Mycobacterium Tuberculosis	1:256
----------------------------	-------

VAPOGENE removes mildew stains from hard surfaces.

### A.O.A.C. OFFICIAL FUNGICIDAL TEST, 11th EDITION

Trichophyton interdigitale	1:200
----------------------------	-------

