

VAPOR SPRAY is a concentrated insecticide for use in control of flying, crawling and running stages of insect life.

Especially compounded for use by food processing establishments such as cameries, wineries, bakeries, restaurants, dairies. Kills accessible stages of flies, mosquitoes, gnats, ants, roaches, silverfish, crickets, spiders, scurpions, wasps, hornets, flying moths, skippers, and most common stored grain pests, including granary weevil, rice weevil, confused flour beetle, cadelle, meal worms, grain moths, grain mites and flour moths.

DIRECTIONS

SURFACE TREATMENT — For control of roaches, silverfish, ants, spiders, exposed stages of skippers and confused flour beetle, granary weevil, rice weevil, cadelle, meai worms, grain moths, grain mites, flour moths and crickets, use good machine adjusted to deliver coarse wet particles. Direct spray onto exposed surfaces of

HIS-14

G-71

VAPOR SPRAY

Pyrethrins 0.30%, Technical Piperonyl butoxide* 1.50%, Petroleum hydrocarbons 98-20%

* Equivalent to 1.20 \circ (butylearbit, \circ (6.3 rops); period (0.3) its 0.30 \circ of related compounds

USDA Reg. No. 3635 141

CAUTION

KEEP OUT OF REACH OF CHILDREN.


HARMFUL IF SWALLOWED.

DO NOT USE, POUR SPILL OR STORE NEAR HEAT OR OPFN FLAME

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when plant is not in operation.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

NET CONTENTS GALLONS



DIRECTIONS CONTINUED

walls, floors, baseboards, cupboards, shelving. Use at rate of 1 gallon per each 1000 sq. ft. of surface area. Apply liberally, paying particular attention to cracks, crevices, inside of empty grain bins and other hiding places. Do not fill grain bins until they are dry after spraying.

SPACE SPRAYING For rapid control of fees mosquitoes, gnatwasps, hornets and flying moths, use in a machine adjusted a deliver a fine mist. Direct spray towards upper portions of the room. Fill the room with mist using 1 to 2 oz per each 1000 cu. ft or space. Where possible, keep room closed for 15 minutes after spraying.

Insecticide use rates vary with different types of thermal for generators, but most units are designed to give most effective results when this material is used at the rate of 1 pint per each 12,000 to 14,000 cu. ft. of space. May be used around food and food products but should not be applied directly onto uncovered foods. Do not remain in treated areas and ventilate the area after treatment is completed.

MANUFACTURED BY



P.O. BOX 80202 ATLANTA, GEORGIA 30341