

Supercide Brand

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes.

Use the product in food processing plants only when plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned with detergent and water before using. Do not spray on food. Thoroughly ventilate treated area before re-entry.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS

Extinguish open flame before space spraying. DO NOT USE. POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

1. Flies and Mosquitoes: Close doors and windows. Direct spray towards ceiling and fill entire room with fog. Keep room closed for 10 to 20 minutes. Do not remain in treated areas and ventilate the area after treatment is completed. Sweep up and destroy fallen insects. Repeat treatment as frequently as required.

I.C.C.: Insecticide, O.T.A.

ACTIVE INGREDIENTS

1. Isoparathinic Petroleum Hydrocarbons	99.7174%
2. Piperonyl Butoxide Technical*	0.2004%
3. Pyrethrin I and II	0.0822%
TOTAL	100.0004%

* (Butylcarbyl) (6-propylpiperonyl) ether and related compounds

This product contains not more than 1% synergists.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

If swallowed, do not induce vomiting. Call physician immediately.


See left panel for additional precautionary statements.

EPA Reg. No. 3635-155
EPA Est. 3635-GA-1

Only For Sale To, Use And Storage By
Servicepersons.

NET CONTENTS _____ GALS.

QJ71
ISC-3
(L85)



2. Cockroaches, Waterbugs and Silverfish Thorough and repeated spraying of infested area should be done, hitting as many of the insects as possible. Pay particular attention to spraying in cracks, crevices, baseboards and other hiding places.

3. Fleas Infesting Premises Can be quickly and effectively controlled by spraying floors and cracks of the infested areas. Fleas lurk in carpets and upholstered furniture. Spray liberally in these areas, as well as in baseboards, wall cracks and sleeping quarters of animals for direct contact with these pests. Repeat applications as necessary for control.

4. Control of Horn and Stable Flies For space spraying apply in a mist with animals on the premises. To temporarily protect cows and horses from fly annoyance, spray from a distance of 6 to 8 feet, using 1 to 2 ounces per animal daily as necessary. May be applied to cattle prior to slaughter. Cows should be sprayed before milking, but care should be taken not to spray directly on udders. Also udders should be washed with a sanitizing solution after spraying and before starting to milk.

5. Weevils, Cadelle and Confused Flour Beetles, Flour Moths in Exposed or Accessible Stages Floors, walls and partitions in flour mills, feed mills, food processing plants, empty grain bins and boxcars should be sprayed with enough solution to wet the surfaces, paying attention to cracks and crevices where insects may be harbored. Spray old sacked grain heavily to reduce infestation. In addition, direct a light spray to hit all exposed ends, sides and tops of storage bags. When reinfestation occurs, repeat the application. Watch for migration periods immediately after or during hot weather. This will materially aid in controlling these pests.

6. Spiders Spray directly onto visible spiders and webs with as wet a spray as possible.

DESCRIPTION

For use in Dairies, Meat Packing Plants, Bakeries, Dairy Barns, Flour Mills, Food Processing Plants, Granaries, Warehouses, Boxcars, Stables, Poultry Plants, Restaurants, Super Markets, Bottling Plants — wherever food products are made, stored, shipped or handled. Recommended for use in Hotels, Motels, Institutions, Theaters, Hospitals, Transportation Facilities, Trains, Buses and Planes.

Kills flies, mosquitoes, roaches, waterbugs, fleas, silverfish, exposed stages of flour weevils, grain beetles and meal moths and spiders that come in contact with the product. Do not apply in food processing areas when these areas are in operation. Remove or cover exposed foodstuffs while spraying. Avoid direct spraying of foods. After spraying, thoroughly wash all surfaces on which spray may have deposited with a detergent solution and rinse with potable water. Modern insect control requires regular treatment and good housekeeping.