Staphylocidal — Fungicidal Tuberculocidal Pseudomonacidal - Salmonellacidal

OXFORD*1922 DISINFECTANT - DETERGENT

ACTIVE INGREDIENTS:

Isopropanol, 11.12%; Tetrasodium ethylenediamine tetraacetic acid, 4.00%; Sodium o-benzyi-p-chlorophenate, 3.75%; Sodium para-tertiary-butylphenate, 3.28%; Sodium xylene sulfonate, 3.20%; Sodium lauryl sulfonate, 3.00%; Sodium 4-chloro-2-cyclopentylphenate, 2.35%; Sodium 2,4,5-trichlorophenate, 2.32%.

.66.98%

DESCRIPTION

1922 is recommended for use in hospitals, nursing homes, food processing plants, animal and poultry houses, hatcheries, and veterinary animal hospitals. At 1 ounce per gallon of water, 1922 can be used to disinfect equipment, walls, floors and any hard inanimate surface not harmed by water.

In the ACAC Use Dilution Method, 11th edition, Oxford 1922 will kill the following organisms at 1:128 dilution (1 ounce per gallon of water).

Staphylococcus aureus, Sal. pullorum, Proteus vulgaris, Salmonella choleraesuis, Sal. typhimurium, Streptococcus mitis, Pseudomonas aeruginosa (PRD 10), Sal. typhosa; Escherichia coli.

AOAC Official Fungicidal Test, 11th Edition, Trichophyton interdigitale 1:128. AOAC Tuberculocidal test, Mycobac erium suberculosis T E

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive: Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of foc J. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

FIRST AID In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed. Rinse empty container thoroughly with water. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

EPA Reg. No. 3635-197 EPA Est 3635-GA-1

Net Contents

Gallons PHD-22

TM.

DIRECTIONS

- 1. For disinfecting, cleaning and deodorizing in hospitals and food processing plants and equipment, use a solution of 1 ounce per gallon (1:128). Mop, wipe, or swab this solution on surfaces. Leave in contact 10 minutes. All surfaces coming in contact with food must be thoroughly scrubbed with detergent and rinse with potable water before reuse.
- 2. For disinfecting and cleaning garbage and trash receptacles, sick room equipment, floors, walls, patient rooms, use 1 ounce to 1 gallon of water (1:128) dilution. For porous surfaces, increase concentration to 2 ounces per gallon of water.
- 3. For non-food disinfection in animal and poultry houses, hatcheries, stockyards, zoos, veterinary animal hospitals: Do not use in milking stalls, milk parlors or milk houses. Remove all poultry, animals and feeds from the premises. Remove all litter and manure from floors and other surfaces; empty all water troughs and feed racks. Saturate all surfaces with a solution of 1 ounce Oxford 1922 per gallon of water (1:128). Ventilate buildings and other closed spaces. Do not house poultry, livestock or use equipment until treatment has been absorbed or dried. All treated feeders, fountains and waterers must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.
- 4. Oxford 1922 kills athlete's foot fungi on environmental surfaces. Prepare a solution of 1 ounce per gallon of water (1:128); mop or swab this solution onto floors and other surfaces normally contacted by the feet in locker rooms, shower stalls, bathrooms and around swimming pools.

UNDER 1988 LEADER INSECTICIDE FUNCIONE : DE MODIL MOIDE ACT

3635-197





PO BOX 80202, ATLANTA GLORGIA 30341