

Center Panel  
Steramine Mark I--Hearns  
July 12, 1967

MILKHOUSE BRAND

STERAMINE  
MARK I  
UDDER WASH  
AND  
SANITIZER

ACTIVE INGREDIENTS:

alkyl (50% C<sub>12</sub>, 50% C<sub>14</sub>) 1.7% C<sub>12</sub>  
dimethyl benzyl ammonium chloride  
isopropyl alcohol 1.0%

\*INERT INGREDIENTS: Water, 97.3%

\*Inert Ingredients: Water, 97.3%

CAUTION - 14 pt. type

KEEP OUT OF REACH OF CHILDREN. Harmful if  
in contact with skin. Avoid getting into the eyes  
or on the skin. In case of contact, flush  
with plenty of water. If irritation per-  
sists, get medical attention. Dilute with  
water as directed. Do not mix with soap.  
Do not use in cleaning instead of  
soap. Rinse with water before sani-  
tizing. Do not allow to come in contact  
with food. Do not reuse container.  
Do not use when empty.

- 13 pt. type

Camoraman's Note  
Euer Copy

Left Panel  
Steramine Mark I--Stearns  
July 11, 1967

INSTRUCTIONS FOR USE:

BULK TANK CLEANING

1. The bulk-up tank truck driver should arrive at the tank at least 15 minutes before pickup with tepid water.
2. Make certain the compressor is working.
3. Remove all detachable parts with the exception of the nozzle and the nozzle label. Brush the interior, nozzle and nozzle label thoroughly.
4. Twice each week clean with a solution of 100 parts of water to 1 part of 45° P for 15 minutes.
5. Reassemble and thoroughly rinse with water.
6. Rinse with a solution of STERAMINE Mark I of water (2 parts of STERAMINE Mark I to 100 parts of water) at a temperature of 100° F. (40° C) for 15 minutes. See recommendations.
7. Before use, rinse with tepid water.

BLENDER WASHING

1. Blender and parts should be washed with water and dried with compressed air.
2. A new solution should be used for each day.

GENERAL INFORMATION

When used as a disinfectant, the solution should be applied to the surface to be disinfected. The solution should be applied to the surface to be disinfected. The solution should be applied to the surface to be disinfected.

STERAMINE MARK I BLENDER WASHING INSTRUCTIONS: The solution should be applied to the surface to be disinfected. The solution should be applied to the surface to be disinfected. The solution should be applied to the surface to be disinfected.

**Steramine Mark I - Stearns**  
July 12, 1967

**INSTRUCTIONS FOR UTENSILS & OTHER DAIRY EQUIPMENT**

1. Immediately following use, wash with warm water.
2. Replenish solution of SPANBOND MARK I at the rate of 2 oz. per gallon of water at 160° F. and circulate for 15-30 minutes.
3. Twice each week follow the same procedure with a solution of SPANBOND at the rate of 1 oz. to 2 gal. of water at 140° F. for 20 minutes for the control of scale.
4. Rinse thoroughly with water at 105° F.
5. Sanitize all utensils with STERAMINE MARK I at the rate of 1 oz. to 4 gal. of water (200 PPM) at a temperature of 145° F. Follow local health recommendations. Do not rinse.
6. Before reuse rinse with potable water.

**PIPELINE MILKERS, MILKING MACHINES & PIPELINES**

1. Immediately after milking, circulate system with warm water 110° F until the water runs clear.
2. Circulate a concentration of SPANBOND MARK I at the rate of 2 oz. per gallon of water 160° F for 15-30 minutes opening and draining all valves at this time. Hand brush all parts not coming in contact with the solution.
3. Twice each week circulate a solution of SPANBOND at the rate of 1 oz. to 2 gal. of water at 140° F for 20 minutes to remove scale and thereby reduce the possibilities of a high bacteria count in milkster production.
4. Rinse with warm water 110° F until the water runs clear.
5. Sanitize with a solution of STERAMINE MARK I at the rate of 1 oz. to 4 gal. of water (200 PPM) and circulate according to local health recommendations.
6. Before reuse rinse with potable water.

