

A WARFARIN Rodenticide

R A C K

R A TTM

W I T H

PROLIN[®]

RAT and MOUSE KILLER

ACTIVE INGREDIENTS:	
Warfarin 3-(a-acetylbenzyl)-4- hydroxycoumarin	0.025%
N ¹ -(2-quinoxaliny) sulfanilamide (sulfaquinoxaline)	0.025%
INERT INGREDIENTS:	99.950%
TOTAL:	<u>100.000%</u>

NET CONTENTS 7 LBS.

ACCEPTED

AUG 24 1971

Packed for
STEARNS CHEMICAL CORPORATION
4200 Sycamore Avenue
P. O. Box 3216
Madison, Wisconsin 53704

[®]Trade Mark of Wisconsin Alumni
Research Foundation



Packed for
STEARNS CHEMICAL CORPORATION
4200 Sycamore Avenue
P. O. Box 3216
Madison, Wisconsin 53704

[®]Trade Mark of Wisconsin Alumni
Research Foundation

CAUTION—KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS IN RIGHT PANEL

A WARFARIN Rodenticide

RACK RAT

PROLIN contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produces Vitamin K, an antidote for Warfarin.

PROLIN is a registered trademark of the Wisconsin Alumni Research Foundation for a rodenticide which is manufactured under license from the Wisconsin Alumni Research Foundation.

FOR CONTROLLING RATS

Place 4 to 6 ounces of PROLIN PELLETS in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feedings have stopped. Replace stale, damp or moldy baits with fresh bait.

FOR CONTROLLING MICE

Place tablespoonful amounts of PROLIN PELLETS in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

CAUTION

Baits should be placed in areas inaccessible to children, pets and domestic animals or in tamper proof bait boxes. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

NOTICE

Seller makes no warranty of any kind expressed or implied concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

PROLIN

RAT and MOUSE KILLER


ACTIVE INGREDIENTS

Warfarin (bishydroxycoumarin) 0.05%

Chlorobutol 0.05%

Other inactive ingredients 94.90%

NET WEIGHT 1.00 LBS.



KEEP OUT OF REACH OF CHILDREN