

DIRECTIONS

PIPELINE MILKERS, MILKING MACHINE & PIPELINE

1. Immediately after milking, flush entire system with warm water 110° F until the water runs clear.
2. Circulate a concentration of AMI-CHEM PIPELINE CLEANER at the rate of 1 ounce per 1.2 gallons of water (100 PPM) at 160° F for 20-30 minutes opening and draining all valves at this time. Hand brush all parts not coming in contact with the solution.

Manufactured by:

STEARNS CHEMICAL CORPORATION
 4200 Sylamore Avenue
 Madison, Wisconsin

NET WT. 10 LBS.

AMI-CHEM

PIPELINE CLEANER

CAUTION:

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Trichloro-S-triazinetriene 1.73%
 (Trichloroisocyanuric Acid)
 (available chlorine 1.56%)

Sodium Metasilicate 20.00%
 (anhydrous)

Total 21.73%

INACTIVE INGREDIENTS 78.27%

100.00%

DIRECTIONS

3. Twice each week circulate a solution of STONE-SOLV at the rate of 1 ounce to 2 gallons of water at 145° F for 20 minutes to remove milkstone and cut down the possibilities of a high bacteria count in milkstone formation.
4. Rinse with warm water 110° F until the water runs out clear.
5. Before reuse, sanitize with a solution of AMI-CHEM PIPELINE CLEANER at the rate of 1 ounce to 1.2 gallons of water (100 PPM) and circulate 5-10 minutes or follow local health recommendations.

USDA Reg. # 3640-53

Packed for:
 Associated Merchants, Inc.
 Beaver Dam, Wisconsin

3640-53

2010-53

DIRECTIONS

PIPELINE MILKERS, MILKING MACHINE & PIPELINE

1. Immediately after milking, flush entire system with warm water 110° F until the water runs clear.
2. Circulate a concentration of AMI-CHEM PIPELINE CLEANER at the rate of 1 ounce per 1.2 gallons of water (100 PPM) at 160° F for 20-30 minutes opening and draining all valves at this time. Hand brush all parts not coming in contact with the solution.

Manufactured by:

STEARNS CHEMICAL CORPORATION
4200 Sycamore Avenue
Madison, Wisconsin

NET WT. 10 LBS.

AMI-CHEM

PIPELINE CLEANER

CAUTION:

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Trichloro-S-triazinetriene 1.73%
(Trichloroisocyanuric Acid)
(available chlorine 1.56%)

Sodium Metasilicate 20.00%
(anhydrous)

Total 21.73%

INACTIVE INGREDIENTS 78.27%

100.00%

BEST DOCUMENT AVAILABLE

DIRECTIONS

3

3. Twice each week circulate a solution of STONE-SOLV at the rate of 1 ounce to 2 gallons of water at 145° F for 20 minutes: to remove milkstone and cut down the possibilities of a high bacteria count in milkstone formation.
4. Rinse with warm water 110° F until the water runs out clear.
5. Before reuse, sanitize with a solution of AMI-CHEM PIPELINE CLEANER at the rate of 1 ounce to 1.2 gallons of water (100 PPM) and circulate 5-10 minutes or follow local health recommendations.

USDA Reg. # 3640-53

Packed for:
Associated Merchants, Inc.
Beaver Dam, Wisconsin

3640-53
3640 53