

ACCEPTED
with COMMENTS
in EPA Letter # 2-11-78

NOV 22 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

3640-75

USE DIRECTIONS

GENERAL FARM USES:

1. Cows and Poultry, Houses
and Houses and Equipment
2. INFECTING SOLUTION

3. Dilute MARK 13 to 10 gallons of water (470 PPM active quaternary) to remove dirt, filth and feces from premises, cars, trucks, camp trailers, trailers, etc. after and droppings from floors, walls, surfaces of barns, etc.

4. If steps, troughs or equipment are soiled by poultry, apply MARK 13 to troughs, racks and other feeding and watering appliances, wash with detergent and rinse thoroughly.

5. To disinfect surfaces with MARK 13, use dilution using a brush, cloth, or sponge on floors, sidewalks and steps.

6. Disinfect buildings, barns, boats, camps, and other closed spaces. Do not enter a poultry or employ equipment until treatment has been absorbed and dried.

7. All treated feed racks, mangers, troughs, automatic feeders, fountains, etc. must be thoroughly rinsed with potable water prior to use.

8. To sanitize eggs intended for hatching, wash to remove gross contamination and then immerse briefly in a sanitizing solution prepared by adding one ounce of MARK 13 to eight gallons of water.

9. To disinfect hatching equipment, remove gross contamination, clean with detergent and rinse with water. Prepare disinfecting solution by adding one ounce of MARK 13 to ten gallons of water and apply to surfaces by brush, mop, brush or cloth.

AIRY

CLEANING BULK MILK CANS

As soon as possible after milk has been removed from bulk tank, rinse tank completely with cold water and allow to drain. Place eight gallons of water (at 120° F) in tank and add one ounce (2 tablespoons) MARK 13 to the water in the tank. Scrub tank, mixers and covers and then allow to drain.

SANITIZING MILK EQUIPMENT

Previously cleaned milking equipment, pails, inflations and tubing should be sanitized prior to milking operations by preparing 200 ppm solution of

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WILLOW CREEK

MARK 13

DISINFECTANT • SANITIZER • DEODORIZER

A powerful germicide for use in Poultry, and Swine areas, in Dairy and for general Farm and Home use. Fast Acting. Effective in Hard Water.

ACTIVE INGREDIENTS

n-Allyl (80% Cl), 30% Cl, 5% Cl, 5% Cl dimethyl benzyl ammonium chloride 10.0%
n-Allyl (80% Cl), 32% Cl, dimethyl ethylbenzyl ammonium chloride 10.0%
Ethyl Alcohol 76.0%

INERT INGREDIENTS

DANGER

KEEP OUT OF REACH OF CHILDREN

See right side panel for risks in pre-occupancy and first aid statements. Guaranteed Minimum Efficacy Coefficients by A.C.A. Test Method at 10°C: *Staphylococcus Aureus*, *Salmonella Typhosa*

Non Flammable, Non Volatile, Odorless, Non Corrosive to inanimate surfaces in use dilution. RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD.

EPA Reg. No. 3640-75

EPA Est. No. 3640 WI 1

Net Contents: 1 Gallon

active quaternaries by nine gallons of water. Scrub or prior to beginning of milk

HARD WATER TOLERANCE

This product fulfills the Milk Ordinance (200 ppm) waters up to 600 ppm of tested by the A.C.A. Test. Use all your milk in your local program.

HOMES, RESTAURANTS, FOOD PROCESSING PLANTS

To sanitize previously silverware, apply a sanitizing water using a cloth for

To disinfect and food and garbage pails apply a gallons of water, using a brush or cloth. Do not DANGER

KEEP OUT OF REACH and skin irritation. Do not face shield and rubber gloves. Avoid contamination of food.

FIRST AID

In case of contact, immerse in water for at least 15 minutes. For eyes, flush with water before reuse.

If swallowed, drink plenty of water. If these are not alcohol. Call a physician.

NOTE TO PHYSICIAN

Pre-treat mucous membranes against circulation may be needed.