

1-6-35  
3625



# STOCK FLY KILL

NON-INJURIOUS WHEN USED AS DIRECTED

## DIRECTIONS FOR USE

Spray Ems Stock Fly Kill liberally so that the insects are enveloped in a heavy mist. In cool weather when flies are at rest on walls and ceilings, spray them directly. Repeat spray treatment as often as necessary.

To protect dairy cattle temporarily from the attack of horn flies, stable flies and mosquitoes, treat twice daily with a light application of spray. Envelop the animal in a mist of Ems Stock Fly Kill, applying it from a distance of two to three feet permitting the spray particles to collect upon the coat. Spray 15 or 20 minutes before milking, washing udder and teats with soap and warm water before milking.

DO NOT SPRAY INTO MILK OR MILK UTENSILS.

CAUTION: Horses, mules, hogs may also be sprayed. Under no circumstances should animals be heavily wetted down. Apply 2 1/2 oz per mature animal and work into hide with brush.

For lice on cattle and hogs - spray every other day. Do not apply more than once every 3 weeks.

For control of chicken mites - Spray thoroughly as often as necessary. Kills lice, ticks, etc. including roost supports, nests, dropping boards and other places. Pay special attention to cracks and crevices.

## ACTIVE INGREDIENTS:

|   |     |
|---|-----|
| B'utoxy B'thiocyno dichyl ether                                       | 1%  |
| B'thiocyno ethyl ester of aliphatic acids averaging 10-18 carbon atom | 3%  |
| Petroleum distillate  | 96% |

DISTRIBUTED BY