

CAUTION: Keep out of Reach of Children
(See right side panel for additional cautions)

E. W. SMITH CHEMICAL CO.
15020 E. Proctor Ave. EDgewood 3-5205
City of Industry, Calif. U.S.A. 91746

LEFT PANEL OF LABEL

- DIRECTIONS -

For protection against hornflies, houseflies, stableflies, spray once or twice daily (caution: do not apply more than two ounces per animal at each mist application for fly protection, using 1 to 2 ounces per head applied as a light mist just prior to milking). Note: sprays used for flies will inhibit the development of blood-sucking and biting lice, using about 2.5 ounces on an animal the size of a mature cow and then brushing it into the hair thoroughly so as to reach the hide, paying particular attention to affected areas. This heavier spraying should be repeated only when necessary and not oftener than once in 3 weeks.

Use Ems PYRENONE FLY SPRAY in stable, barns, houses, food processing plants and milk parlors for effective control of flies. Direct the mist toward the upper portions of the enclosure, filling the air with spray. Whenever possible, keep doors and windows closed for 10 minutes after application. Do not remain in treated areas in enclosed areas and ventilate the area after treatment is completed.

RIGHT PANEL OF LABEL

Ems PYRENONE FLY SPRAY is a special insecticide formulation compounded for the livestock, dairy and feed processing industries. It contains Piperonyl butoxide. It is especially effective for rapid and economical control of hornflies. Avoid wetting the hides of animals.

CAUTION: DO NOT STORE, SPRAY OR OTHERWISE EXPOSE TO OPEN FLAME. HARMFUL IF SWALLOWED. Use the product as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

ACTIVE INGREDIENTS:

Piperonyl butoxide technical *	0.60%
Pyrethrine	0.06%
Petroleum Hydrocarbons	99.34%

* Equivalent to 0.48% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.12% of related compounds.

* Reg. trademark FMC

U.S.D.A. Registration No. 3682-28

[3682-28]

H. W. SMITH CHEMICAL CO.
LA PUENTE, CALIF.

FRONT PANEL OF LABEL:

Ems PYRENONE FLY SPRAY

CAUTION: Keep out of Reach of Children
(See right side panel for additional cautions)

E. W. SMITH CHEMICAL CO.
15020 E. Proctor Ave. Edgewood 3-5205
City of Industry, Calif. U.S.A. 91746

LEFT PANEL OF LABEL

- DIRECTIONS -

For protection against hornflies, houseflies, stableflies, spray once or twice daily (caution: do not apply more than two ounces per animal at each mist application for fly protection, using 1 to 2 ounces per head applied as a light mist just prior to milking). Note: sprays used for flies will inhibit the development of blood-sucking and biting lice, using about 2.5 ounces on an animal the size of a mature cow and then brushing it into the hair thoroughly so as to reach the hide, paying particular attention to affected areas. This heavier spraying should be repeated only when necessary and not oftener than once in 3 weeks.

Use Ems PYRENONE FLY SPRAY in stable, barns, houses, food processing plants and milk parlors for effective control of flies. Direct the mist toward upper portions of the enclosure, filling the air with spray. Whenever possible, keep doors and windows closed for 10 minutes after application. Do not remain in treated areas in enclosed areas and ventilate the area after treatment is completed.

RIGHT PANEL OF LABEL

Ems PYRENONE FLY SPRAY is a special insecticide formulation compounded for the livestock, dairy and feed processing industries. It contains Piperonyl butoxide. It is especially effective for rapid and economical control of hornflies. Avoid wetting the hides of animals.

CAUTION: DO NOT STORE, SPRAY OR OTHERWISE EXPOSE TO AN OPEN FLAME. HARMFUL IF SWALLOWED. Use the product as a space spray in food processing plants when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

ACTIVE INGREDIENTS:

Piperonyl butoxide technical *	0.60%
Pyrethrine	0.06%
Petroleum Hydrocarbons	99.34%

* Equivalent to 0.48% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.12% of related compounds.

* Reg. trademark FMC

U.S.D.A. Registration No. 3682-28