

FEB 23, 1973

Kingstree, S. C. 29556

4% MALATHION DUST

## WITH SULPHUR

ACTIVE INGREDIENTS	BY WEIGHT	Γ
Malathion	49	6
Sulphur	189	ć
INERT INGREDIENTS	78%	ć
TOTAL	100;	ć

\*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

## CAUTION: KEEP OUT OF REACH OF CHILDREN

BEANS: For control of Mexican Bean Beetle, Bean Leaf Hopper, Spider Mites, Thrips and Aphids, apply at the rate of 30-35 pounds per acre. Begin applications early and continue every 7-10 days. For best results apply to underside of leaves. Do not apply to beans within 3 days of harvest.

CABBAGE: For control of Aphids and imported Cabbageworms, apply at the rate of 30-35 pounds per acre. Begin applications when aphids or worms first appar and continue every 7-10 days. Do not apply within 7 days of harvest.

CAUTION: Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and food-stuffs.

Remove harmful pesticidal residues at harvest from edible parts of growing food crops by washing, brushing or other means.

## DO NOT REUSE EMPTY BAG. BURY WITH WASTE IN A SAFE PLACE.—

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

E.P.A. REG. NO. 3743-64