





DANGER--POISON--PRECAUTIONS

POISONOUS IF SWALLOWED

This product can kill you if swallowed. Even small amounts can be fatal. Do not swallow.

POISONOUS BY SKIN CONTACT

Product absorbed by skin is irritating. Washed hands thoroughly with soap and water. Do not touch face through clothes.

POISONOUS IF BREATHED

Irritating vapors, spray mist or dust may be fatal.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep off any body of water. Do not apply when weather is likely to favor drift from treated areas. Do not use near water freely flowing.

POISONOUS TO FISH

POISONOUS

POISONOUS

POISONOUS

STOP READ LABEL BEFORE USING!

POISON

See side panels
for antidote
and precautions



DANGER

Keep out of reach
of children

CAN KILL YOU

PARATHION - 4C

NOT FOR HOME USE

Southern Agricultural Chemicals Co.
KING, TREE, & NEWLAND

Apply 2-4 weeks before planting into the upper 4 inches of soil. Do not use in home garden.

STOP - READ LABEL BEFORE USING!

POISON

See side panels
for antidote
and precautions



DANGER

Keep out of reach
of children

CAN KILL YOU

PARATHION-4G

NOT FOR HOME USE

NET WEIGHT

MANUFACTURED BY

Southern Agricultural Chemicals, Inc.
KINGSTREE, S. C. NEWLAND, N. C.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

(1) Wear water proof pants, coat, hat, rubber boots or rubber over shoes. (2) Wear safety goggles. (3) Wear mask or respirator approved by the U.S. Government. Masks for parathion protection. (4) Wear heavy duty, natural rubber gloves.

WORK SAFETY RULES

Keep all unprotected persons and children away from treated areas where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick, stop work immediately and get away from a doctor, physician, nurse, or hospital. Explain that the victim has been exposed to parathion and describe the symptoms. Go to the nearest hospital. See First Aid Treatment Section and First Aid Manual for more information.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before sleeping, eating, or drinking.

AFTER WORK, take off all work clothes and shoes. Shower with soap and water. Wear clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

POISON SIGNS (Symptoms)

Parathion is a very dangerous poison. It rapidly enters the body on contact with all body surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment immediately.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, dizziness, weakness, blurred vision, pinpoint pupils, tightness in chest, labored breathing, convulsions, sweating, watering of eyes, drooling or foaming of mouth and nose, loss of vision and coma.

FIRST AID TREATMENT

Get victim to a doctor or hospital immediately. If victim is unconscious, do not give anything by mouth and do not give anything by mouth.

If victim is conscious, start artificial respiration. If victim is unconscious, make him or her breathe.

If victim has swallowed and victim is alive, give 1-2 glasses of water to drink. If victim is unconscious, do not give anything by mouth.

If victim has inhaled, remove victim from exposure. If victim is unconscious, do not give anything by mouth.

If victim has inhaled, remove victim from exposure. If victim is unconscious, do not give anything by mouth.

NOTE TO PHYSICIAN

Parathion is a very dangerous poison. It rapidly enters the body on contact with all body surfaces and eyes. Clothing wet with this material must be removed immediately.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, dizziness, weakness, blurred vision, pinpoint pupils, tightness in chest, labored breathing, convulsions, sweating, watering of eyes, drooling or foaming of mouth and nose, loss of vision and coma.

First Aid Treatment: Get victim to a doctor or hospital immediately. If victim is unconscious, do not give anything by mouth and do not give anything by mouth.

If victim is conscious, start artificial respiration. If victim is unconscious, make him or her breathe. If victim has swallowed and victim is alive, give 1-2 glasses of water to drink.

If victim has inhaled, remove victim from exposure. If victim is unconscious, do not give anything by mouth. If victim has inhaled, remove victim from exposure. If victim is unconscious, do not give anything by mouth.

EPA Reg. No. 371-175