

5% SEVIN* DUST

ACTIVE INGREDIENT:
Carbaryl (1-Naphthyl N-Methylcarbamate)
INERT INGREDIENTS:

By Wt
5.00
95.00%
100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Trade Marks Reg. U. S. Patent Office

*Sevin Trade Mark of UCC for 1-Naphthyl N-Methylcarbamate

5% Sevin Dust is an insecticidal dust for use against certain insects attacking beans and potatoes.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

TOBACCO—For control of Hornworms, Budworms, Flea Beetles, use at rate of 15-20 lbs. per acre. Treat infested fields before largest worms exceed 1 1/2 inches. When treating during harvest make applications immediately after, rather than before priming.

BEANS— Against Mexican Bean Beetle, dust thoroughly with 20 Lbs. per acre, every 7 to 10 days or as needed, beginning when insects first appear. Against Leafhoppers use 30-40 Lbs. per acre.

IRISH POTATOES—Against Colorado Potato Beetle, dust thoroughly using 20 Lbs. per acre. Against Leafhoppers and Flea Beetles use up to 30 Lbs per acre. Dust when insects appear and repeat every 7 to 10 days or as needed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE

The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues. Residue tests indicate that this dust can be applied at rates directed above to beans and potatoes on the day of harvest without exceeding residue tolerances.

CAUTION

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED OR INHALED. Do Not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of food and foodstuffs. Do not feed treated vines to meat or dairy animals.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY

EPA REG. NO. 3743-134

