STREPTOMYCIN ACCEPTED ANTIBIOTIC DUST

A FIELD STRENGTH DUST FOR CONTROL OF CERTAIN FUNGUS AND BACTERIAL DISEASES IN PLANTS.

Transport ACTIVE INGREDIENTS

Streptomycin _____

Imentx incredients _____

98.8%

INERT INC. EDIENTS

DANGER: KEEP OUT OF REACH OF CHILDREN

This material is non-sterile grade of Streptomycin and is not to be used for medicinal purposes.

DIRECTIONS FOR USE:

Wildfire and Blue Mold in TOBACCO PLANT BEDS-

PREVENTIVE TREATMENT—Apply Streptomycin-two by dusting 2 to 3 lbs. evenly over 100 square yards of plant bed when plants are in 2 leaf stage. Apply at least four dustings at 5 to 7 day intervals. Increase dosage to 3 to 4 pounds for last 2 dustings.

CURATIVE TREATMENT-In beds where wildfire or bluemold is already established, dust 4 pounds evenly over 100 yards of plant bed. Repeat two to three times as necessary.

CAUTIONS

May cause skin irritation. Avoid prolonged or repeated contact with skin. In case of contact flush with plenty of water.

NOTICE OF WARRANTY

The manufacturer makes no warranty, express or implied, concerning the use of this product. The user assumes all risks and liability in connection with its use.

Reg. No. 3743-136