


The Sign of Quality



4% MALATHION DUST

ACTIVE INGREDIENTS	BY WEIGHT
Malathion	4%
INERT INGREDIENTS	96%
TOTAL	100%

*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION: KEEP OUT OF REACH OF CHILDREN

BEANS: For control of Mexican Bean Beetle, Bean Leaf Hopper, Spider Mites, Thrips and Aphids, apply at the rate of 30-35 pounds per acre. Begin applications early and continue every 7-10 days. For best results apply to underside of leaves. Do not apply to beans within 3 days of harvest.

CABBAGE: For control of Aphids and imported Cabbageworms, apply at the rate of 30-35 pounds per acre. Begin applications when aphids or worms first appear and continue every 7-10 days. Do not apply within 7 days of harvest.

POULTRY: Lice and Mites in Poultry House, dust 1 lb. per 40 sq. ft. of litter, giving thorough coverage of house, roosts and nests. Replace nesting material after dusting. Do not apply while birds are in house.

CAUTION: Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and food-stuffs.

KEEP OUT OF REACH OF CHILDREN.

This Product will kill fish. Keep out of lakes, streams, and ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Remove harmful pesticidal residues at harvest from edible parts of growing food crops by washing, brushing or other means.

DO NOT RE-USE EMPTY CONTAINER. DESTROY WHEN EMPTY.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

E. P. A. Reg. No. 3743-149