


Small illegible text



CAN KILL YOU

PARATHION ETHIODAN 2-4-4

Manufactured By
Southern Agricultural Chemicals, Inc.
KINGSTREE, VA

AC 2-4-4

(Contains 1.0 lb. of actual Parathion and 2.0 lbs of actual Thiodan per gallon)

DIRECTIONS FOR USE

Apply the dosages indicated per acre in the proper amount of water to give through coverage of the plant.

TOBACCO: To control budworm and tobacco suckly, use 1½ to 2 pints per acre. To control cabbage looper, flea beetle, Green Peach Aphid and hornworm, use 1 to 2 quarts per acre. For vegetable weevil, use 2 quarts per acre. Note: Do not apply within 5 days of priming or within 15 days of cutting. Do not enter fields within 24 hours of application. Protective clothing should be worn during applications.

TOBACCO (PLANT BED TREATMENT): Green June Bug Larvae, 1 pint per 100 square feet of plant bed. mixture per square yard of plant bed.

TOMATOES: To control aphids, blister beetle, colorado potato beetle, flea beetle, green stink bug, whitefly use 1 quart per acre. Note: Do not apply within 10 days of picking. To control Hornworm use 1 to 2 quarts per acre. Note: Do not apply within 10 days.

CUCUMBERS, PUMPKINS, SUMMER SQUASH, WINTER SQUASH: To control aphids, cucumber beetles use 1 to 2 quarts per acre. Note: Do not apply before vining or within 15 days of harvest.

COTTON: To control spider mites, (desert spider mite, tumid spider mite, two-spotted spider mite), aphids, cotton leafworm, lygus bug, bollworm, cabbage looper use 2 quarts per acre. Note: Do not apply after bolls begin to open. Do not graze dairy or meat animals in treated fields.

POTATOES: To control flea beetle, colorado potato beetle, leafhopper, aphids, southern armyworm, green stink bug, leaf-footed plant bugs and other plant bugs, potato tuberworm, whitefly, use 1 to 2 quarts per acre. Note: Do not apply within 5 day of harvest. To control false chinch bug use 2 quarts per acre. Note: Do not apply within 5 days of harvest. To control potato psyllid use 1½ to 2 quarts per acre. Note: Do not apply within 5 days of harvest.

BROCCOLI, CABBAGE: To control cabbage aphids, cabbage looper, imported cabbage-worm, diamond-back moth, larvae, cross-striped cabbageworm, flea beetle, harlequin bug, stink bug, use 1½ to 2 qts. per acre. Note: Do not apply within 10 days of harvest on cabbage or within 7 days of harvest on Broccoli.

To protect fish and wild life, do not contaminate streams, lakes or ponds with this material.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

F.P.A. Reg. No. 3743-155

