

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**THIS PRODUCT IS NOT FOR RESALE
TO BE USED FOR MANUFACTURING PURPOSES ONLY
STOP - READ LABEL BEFORE USING!**



DANGER
Keep out of reach
of children

CAN KILL YOU

PARATHION-25DB

NOT FOR HOME USE

APPENDICES

the 1990s, the number of children with disabilities has increased. The following table shows the number of children with disabilities in the United States in 1990 and 2000.

NET VOLUME

MANUFACTURED BY

Southern Agricultural Chemicals, Inc.
KINGSREE, S. C. NEWLAND, N. C.

GALLON

DANGER—POISON—PRECAUTIONS 



POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts. spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Personnel touched by hands or exposed pushed
on skin in eyes, mouth, clothing, body, through
clothes)

POISONOUS IF BREATHED

Revised: 10/15/2015

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed, being out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding the best management practices.

CONTAINER DISPOSAL

BEST DOCUMENT AVAILABLE

USE ONLY WHEN WEARING THE FOLLOWING CLOTHING

(1) Wear water proof pants, coat, hat, rubber boots or goggles. (3) Wear mask or respirator approved by the U.S. protection. (4) Wear heavy duty, natural rubber gloves.

WORK SAFETY RULE

Keep all unprotected persons and children away from
danger of drift

From the eyes or mouth with hands. If you feel sick -
right away. Call a doctor (physician), clinic or hos-
pital. If you have been exposed to parathion and describe this
free First Aid Treatment Section) and if a doctor can't
help, call hospital.

IMPORTANT: Before removing gloves, wash them w

IMPORTANT: Before removing gloves, wash them with hands, face and arms with soap and water before smoking.
AFTER WORK: Take off all work clothes and shoes. So only clean clothes when leaving job. Do not wear contaminated and protect equipment with soap and water, be cleaned and later replaced according to instructions.

POISON SIGNS (Symptoms)

Parathion is a very dangerous poison. If rapidly enters surfaces and eyes, rinsing well with this material immediately is most useful. The prompt medical treatment of the signs and symptoms of poisoning are the least important thing to do. Do not let up on tightness, loss of sweating, watering of eyes, drooping of the eyelids, and so on.

FIRST AID TREATMENT

[illegible]

NOTE TO PHYSICI

A. The patient is generally responsive to large doses. The patient is also responsive to the 500 mg dose. The patient is also responsive to the 100 mg dose. The patient is also responsive to the 50 mg dose. The patient is also responsive to the 25 mg dose. The patient is also responsive to the 12.5 mg dose. The patient is also responsive to the 6.25 mg dose. The patient is also responsive to the 3.125 mg dose. The patient is also responsive to the 1.5625 mg dose. The patient is also responsive to the 0.78125 mg dose. The patient is also responsive to the 0.390625 mg dose. The patient is also responsive to the 0.1953125 mg dose. The patient is also responsive to the 0.09765625 mg dose. The patient is also responsive to the 0.048828125 mg dose. The patient is also responsive to the 0.0244140625 mg dose. The patient is also responsive to the 0.01220703125 mg dose. The patient is also responsive to the 0.006103515625 mg dose. The patient is also responsive to the 0.0030517578125 mg dose. The patient is also responsive to the 0.00152587890625 mg dose. The patient is also responsive to the 0.000762939453125 mg dose. The patient is also responsive to the 0.0003814697265625 mg dose. The patient is also responsive to the 0.00019073486328125 mg dose. The patient is also responsive to the 0.000095367431640625 mg dose. The patient is also responsive to the 0.0000476837158203125 mg dose. The patient is also responsive to the 0.00002384185791015625 mg dose. The patient is also responsive to the 0.000011920928955078125 mg dose. The patient is also responsive to the 0.0000059604644775390625 mg dose. The patient is also responsive to the 0.00000298023223876953125 mg dose. The patient is also responsive to the 0.000001490116119384765625 mg dose. The patient is also responsive to the 0.0000007450580596923828125 mg dose. The patient is also responsive to the 0.00000037252902984619140625 mg dose. The patient is also responsive to the 0.000000186264514923095703125 mg dose. The patient is also responsive to the 0.0000000931322574615478515625 mg dose. The patient is also responsive to the 0.00000004656612873077392578125 mg dose. The patient is also responsive to the 0.000000023283064365386962890625 mg dose. The patient is also responsive to the 0.0000000116415321826934814453125 mg dose. The patient is also responsive to the 0.00000000582076609134674072265625 mg dose. The patient is also responsive to the 0.000000002910383045673370361328125 mg dose. The patient is also responsive to the 0.0000000014551915228366851806640625 mg dose. The patient is also responsive to the 0.00000000072759576141834259033203125 mg dose. The patient is also responsive to the 0.000000000363797880709171295166015625 mg dose. The patient is also responsive to the 0.0000000001818989403545856475830078125 mg dose. The patient is also responsive to the 0.00000000009094947017729282379150390625 mg dose. The patient is also responsive to the 0.000000000045474735088646411895751953125 mg dose. The patient is also responsive to the 0.0000000000227373675443232059478759765625 mg dose. The patient is also responsive to the 0.00000000001136868377216160297393798828125 mg dose. The patient is also responsive to the 0.000000000005684341886080801486968994140625 mg dose. The patient is also responsive to the 0.0000000000028421709430404007434844970703125 mg dose. The patient is also responsive to the 0.00000000000142108547152020037174224853515625 mg dose. The patient is also responsive to the 0.000000000000710542735760100185871124267578125 mg dose. The patient is also responsive to the 0.0000000000003552713678800500929355621337890625 mg dose. The patient is also responsive to the 0.00000000000017763568394002504646778106689453125 mg dose. The patient is also responsive to the 0.000000000000088817841970012523233890533447265625 mg dose. The patient is also responsive to the 0.0000000000000444089209850062616169452667236328125 mg dose. The patient is also responsive to the 0.00000000000002220446049250313080847263336181640625 mg dose. The patient is also responsive to the 0.000000000000011102230246251565404236316680908203125 mg dose. The patient is also responsive to the 0.00000000000000555111512312578270211815833404940625 mg dose. The patient is also responsive to the 0.000000000000002775557561562891351059079167024703125 mg dose. The patient is also responsive to the 0.0000000000000013877787807814456755295395835123515625 mg dose. The patient is also responsive to the 0.00000000000000069388939039072283776476979175617578125 mg dose. The patient is also responsive to the 0.000000000000000346944695195361418882384895878087890625 mg dose. The patient is also responsive to the 0.0000000000000001734723475976807094411924479390439453125 mg dose. The patient is also responsive to the 0.00000000000000008673617379884035472059622396952197265625 mg dose. The patient is also responsive to the 0.000000000000000043368086899420177360298111984760986328125 mg dose. The patient is also responsive to the 0.0000000000000000216840434497100886801490559923804931640625 mg dose. The patient is also responsive to the 0.00000000000000001084202172485504434007452799619024658203125 mg dose. The patient is also responsive to the 0.000000000000000005421010862427522170037263998095123291015625 mg dose. The patient is also responsive to the 0.000000000000000002710505431213761085018631999047561645578125 mg dose. The patient is also responsive to the 0.0000000000000000013552527156068805425093159995237808227890625 mg dose. The patient is also responsive to the 0.00000000000000000067762635780344027125465799976189041139453125 mg dose. The patient is also responsive to the 0.000000000000000000338813178901720135627328999880945205697265625 mg dose. The patient is also responsive to the 0.0000000000000000001694065894508600678136644999404726028486328125 mg dose. The patient is also responsive to the 0.00000000000000000008470329472543003390683224997023630142431640625 mg dose. The patient is also responsive to the 0.000000000000000000042351647362715016953416124985118150712158203125 mg dose. The patient is also responsive to the 0.0000000000000000000211758236813575084767080624925590753560791015625 mg dose. The patient is also responsive to the 0.0000000000000000000105879118406787542383540312462795376780395578125 mg dose. The patient is also responsive to the 0.00000000000000000000529395592033937711917701562313976883901977890625 mg dose. The patient is also responsive to the 0.000000000000000000002646977960169688559588507811698944419509889453125 mg dose. The patient is also responsive to the 0.0000000000000000000013234889800848442779792539055849722097549447265625 mg dose. The patient is also responsive to the 0.00000000000000000000066174449004242213898962695279248610487747236328125 mg dose. The patient is also responsive to the 0.000000000000000000000330872245021211069494813476396243052438736181640625 mg dose. The patient is also responsive to the 0.0000000000000000000001654361225106055347474067381981215262193680908203125 mg dose. The patient is also responsive to the 0.00000000000000000000008271806125530276737370336909906076310968404540986328125 mg dose. The patient is also responsive to the 0

Consult your State A
post to transfer assets

— — — — —

BEST DOCUMENT AVAILABLE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

STOP - READ LABEL BEFORE USING!

STOP - READ LABEL BEFORE USING!

STOP - READ LABEL BEFORE USING!

**Keep out of reach
of children**

CAN KILL YOU

PARATHION-25DB

NOT FOR HOME USE

ACTIVE INGREDIENTS

PARAFFIN OIL, 100.00 g. (100.00 ml.)
INERT INGREDIENTS, 10.00 g.

0011A1

1100

NET VOLUME

MANUFACTURED BY

Southern Agricultural Chemicals, Inc.

KINGSTREE, S. C. NEWLAND, S. C.

BEST DOCUMENT AVAILABLE

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

1) Wear water-tight pants, coat, hat, rubber boots or rubber overshoes. (2) Wear safety glasses. (3) Wear mask or respirator approved by the U.S. Bureau of Mines for parathion protection. (4) Wear heavy-duty natural rubber gloves.

WORK SAFETY RULES

Keep a safe distance from persons and children away from treated area or where there is spraying of spray.

Call for help or call 911 with hands if you feel sick in any way. STOP work and get help right away. A sick worker should go to clinic or hospital immediately. Explain that the worker has been exposed to pesticides and describe the condition. After first aid is given, use First Aid treatment sheet and if a doctor cannot come, take victim to clinic or hospital.

IMPORTANT: Do not use a glass with ice with soap and water. Always wash
hands before eating, drinking, and water for drinking.

AFTER WORK: Wash face, work clothes and shoes. Shower using soap and water. Wear protective gloves when handling oil. Do not wear contaminated gloves with no wash protection. Store and protect the respirator with a strap and water after each use. Respirator should be cleaned after each use. Do not use contaminated gloves to clean a respirator.

POISON SIGNS (Symptoms)
Parathion is a very strong poison. If it enters the body on contact with the skin or is inhaled, it can be fatal. If it is swallowed, it must be removed immediately. Ex-
posure to the skin or eyes causes a burning sensation. If it is swallowed, it causes

POISON SIGNS (Symptoms)

[illegible]

FIRST AID TREATMENT

14. A letter dated 10-1-68 from the above named person has been received.

the patient's past history of a fall, a seizure, a recent fall, and a more an unusual accident. The patient was not aware of any other cause for the vomiting. The patient was not aware of any other cause for the vomiting. The patient was not aware of any other cause for the vomiting.

2. The water is not only a source of drinking water for all, but is also a source of water for the people of the world.

NOTE TO PHYSICIAN

[illegible]

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

Application should be made only by experienced personnel using approved safety equipment. Do not use for Aircraft application. Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to settle. If mixture does not mix the tendency to separate, do not use. Injury to plants may result. Bury any unused mixture in a safe place. Do not combine emulsifiable liquid with wettable powder in the same spray tank unless previous use of the material demonstrated no possible injury to plants is possible.

RECOMMENDATIONS

BEANS: Aphids, Armyworms, Leafhoppers, Leaf Miners, Leaf Beetles, and Mexican Bean Beetle. Use 1 pt. per acre in sufficient water for thorough coverage. Repeat applications at 7 to 10 day intervals until harvest. Do not apply within 15 days of harvest.

BEETS: Aphids, Weevils. Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply within 15 days of harvest (21 days if treated tops are to be used for food or feed).
COLE CROPS: Cabbage, Broccoli, Cauliflower, Kohlrabi, Turnips, Turnip Greens, and Rutabaga. Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply within 15 days of harvest.

DESTROY EMPTY CONTAINERS
Perforate, Crush

COTTON: Aphids, Red Spider Mites. Use $\frac{1}{4}$ to 2.5 pt. per acre in sufficient water for thorough coverage. Do not hand pick or harvest within 7 days.

CUCURBITS: (Cucumber, Squash and Melons) Aphids, Cucumber Beetle, Melonworm, Pickworm, Serpentine Leaf-miner. Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply unless plants are dry nor until tomatoes begin to form. Do not apply within 15 days of harvest on cucumbers and squash; 7 days on melons.

FORAGE CROPS: Alfalfa, Barley, Clover, Oats, Vetch, Wheat, Apples, Armyworm, Bollworm, Boll Beetle, Cotton, Eggplant. Use $\frac{1}{4}$ to 2.5 pt. per acre in sufficient water for thorough coverage at first insect infestation. Do not apply within 15 days of harvest or forage use.

POTATOES, TOMATOES: Aphids, Flea Beetle, Leaf miner, Colorado Potato Beetle. Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply within 10 days of harvest or within 15 days of first insect infestation.

NOTICE: This material is highly toxic to fish and many other aquatic life. Do not use in or near water. Do not use in or near streams, rivers, lakes, ponds, or other bodies of water. Do not use in or near areas where water is used for drinking or irrigation. Do not use in or near areas where water is used for fishing or other recreational purposes. Do not use in or near areas where water is used for livestock or poultry consumption. Do not use in or near areas where water is used for human consumption. Do not use in or near areas where water is used for industrial or commercial purposes. Do not use in or near areas where water is used for other purposes.



DANGER—POISON—PRECAUTIONS

POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts. Spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Prolonged contact with skin or clothing splashed with this product may cause irritation and may be fatal if absorbed through the skin.

POISONOUS IF BREATHED

Exposure to vapors may cause irritation and may be fatal.

POISONOUS TO FISH AND WILDLIFE

This product is highly toxic to fish and many other aquatic life. Do not use in or near water. Do not use in or near streams, rivers, lakes, ponds, or other bodies of water. Do not use in or near areas where water is used for drinking or irrigation. Do not use in or near areas where water is used for fishing or other recreational purposes. Do not use in or near areas where water is used for livestock or poultry consumption. Do not use in or near areas where water is used for human consumption. Do not use in or near areas where water is used for industrial or commercial purposes. Do not use in or near areas where water is used for other purposes.

POST TREATED AREA