

CUCUMBER, MELONS, PUMPKINS, SQUASH: Pickleworm, Melonworms — 1 lb. per acre in water to cover. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs — 2 lbs. per acre in water to cover. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use Sevin on Watermelons in Florida.

LAWNS: Chinch Bugs, Leafhopper, Fall Armyworm, Sod Webworms or Lawn Moths — Use 1 cupful in 15 to 20 gals. water for each 500 sq. ft. lawn area. Use garden hose sprayers or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat 2 to 3 weeks later if necessary.

ORNAMENTALS: (Andromeda, Arborvitae, Ash, Aster, Azalea, Bamboo, Beech, Begonia, Birch, Boxwood, Camellia, Carnation, Ceanothus, Cedar, Chinese Elm, Chrysanthemum, Cypress, Dahlia, Delphinium, Dogwood, Elm, Evonymus, Fir, Fuchsia, Gladiolus, Hemlock, Hickory Holly, Hornbeam, Hydrangea, Inkberry, Iris, Juniper, Laurel, Lillac, Locust, Magnolia, Maidenhair fern, Maple, Marigold, Nasturtium, Oak, Orchid, Ornamental, Mulberry, Petunia, Pine, Poplar, Pivet, Rhododendron, Rose, Rose of Sharon, Sassafras, Spirea, Verbena, Viburnum, Willow, Wisteria, Yew, Zinna): Apple, Rose, and Elm Leaf Aphid, Bagworm, Blister Beetles, Boxelder Bug, Certain Leaf Miners, Elm Leaf Beetle, Flea Beetles, Japanese Beetle, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Pland Bugs, Scale Insects, Ten Caterpillars, Exposed Thrips, Lace Bugs, Psyllids, Rose-Slug, Willow Leaf Beetle — 2 lb. per 100 gals. water (2 tablespoonful, per gal.)

Potato, Eggplant Pepper: Colorado Potato Beetle, Tarnished Plant Bug — 1 to 2 lbs. per acre in water to cover. Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworms, Tomato Hornworms, European Corn Borer, Stink Bugs, Lygus Bugs — 2 lbs. per acre in water to cover. May be applied up to the day of harvest.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leaf Roller — 2 lbs. per 100 gals. water (200 gals. per acre). Make full coverage application. Apply in summer sprays. Repeat 3 to 5 times when insect appears. Preharvest Limit — 1 Day.

TOBACCO: (In Plant Beds — Except Kentucky): Tobacco Flea Beetle — apply 2 lbs. per 50 gals. water and apply 6 gals. per 100 sq. yds. when plants appear. Repeat as needed at 10 to 14-day intervals and just before transplanting (In Fields — All Areas) — Hornworms, Tobacco Flea Beetles — Use 2 to 4 lbs. in 50 to 100 gals. water per acre, depending on size of plant. Use higher dosage on larger plants and repeat as needed. Good coverage is essential. Heavier applications may cause plant injury. If late applications are necessary, treat immediately after priming. Do not apply within 5 days of priming or 15 days of cutting. Avoid getting wet with plant juices when cutting the crop.

CABBAGE: Diamondback moth, Imported cabbage worm, Cutworm, and Flea Beetle — 2-4 lbs. per acre in water to cover. Apply on foliage as needed.

TOMATO: Blister Beetle, Colorado potato beetle, Cutworm, Flea Beetle — 2-4 lbs. per acre in water to cover. Hornworm, Stink bug, Tomato fruitworm — 4 lbs. per acre in water to cover. Apply to foliage as needed.

COMPATIBILITY: This product is compatible with commonly used insecticides such as lead arsenate and chlorinated hydrocarbons, Parathion and other organic Phosphates, Botanicals, also with fungicides as Organo-mercury compounds, Fixed Cop-pers, Carbamates, Sulfur, Captain, Cyprax, Ciyodin and Phalton. Do not use with lime, Bordeaux, Lime-sulfur, Casein-Lime spreaders of other alkaline materials.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing of Dust or Spray. Do not take internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physicians: Carbaryl is a moderate reversible, Cholin-esterase Inhibitor. Atrophine is antidotal. **AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.** This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply under conditions involving possible drift to food forage or other plant-ings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Do not reuse Container. Destroy when empty. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MANUFACTURED BY
SOUTHERN AGRICULTURAL CHEMICALS, INC.
Kingstree, S. C.

ROYAL BRAND SEVIN 50% WETTABLE

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-methylcarbamate) 50%

INERT INGREDIENTS

50%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS, AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

CAUTION: Keep Out of Reach of Children

SEE OTHER PRECAUTIONS ON THE BACKSIDE PANEL.

Keep spray mixtures well agitated. Sevin 50 Wettable may be applied within 1 day of harvest on all food crops listed unless otherwise noted.

APPLES, PEARS (WESTERN): Apple Sucker, Green Apple Aphid, Woolly Apple Aphid, Bagworm California, Pear-slug, Codling Moth, Apple Rust Mite, Pear Leaf Blister Mite, Eye Spotted Bud Moth, Green Fruitworm, Lygus Bugs, Orange Tortrix, Pear Rust Mite, Pear Psylla, Tentiform Leaf Minors, Lecanium Scales, Oystershell Schies, San Jose Scale — 1½ to 2 lbs. per 100 gals. water. For Best Scale Control — Apply when crawlers are present. For Psylla — apply when eggs hatch or when young nymphs are present. Heavy Aphid Infestations — Repeat as necessary for best control. (EASTERN STATES): Apple Mealybug, green Apple Aphid, Codling Moth, White Apple Leafhopper — 1 lb. per 100 gals. water. Heavy Aphid Infestations — Repeat as necessary for best control. Apple Maggot Bagworm, Eastern Tent Caterpillar, European Apple sawfly, Eye-spotted Bud Moth, Fruit Tree Leaf Roller, Green Fruitworm, Japanese Beetle, Pear Psylla, Periodical Cicada, Plum Curculio, Red-banded Leaf Roller, Rosy Apple Aphid, Woolly Apple Aphid, Apple Rust Mite, Pear Leaf Blister Mite, Pear Rust Mite, Tarnished Plant Bug, Tentiform Leaf Minors, Forbs Scale, Lecanium Scales Oystershell Scale, San Jose Scale — 2 lbs. per 100 gals. water. For Best Scale Control — Apply when crawlers are present. Rosy Apple Aphid — Apply before leaves are curled. Apple Maggot — Apply at 10-day intervals. To avoid possible fruit thinning, do not apply in any area within 30 days after bloom. Foliar injury may result from combination with summer oils. May cause foliar injury if used before second cover on York and McIntosh apples. Preharvest Limit — 1 day. Apply at Petal Fall and every 10 to 14 days thereafter until control is achieved.

APPLE THINNING: Easily-Thinned Varieties such as Cortland, Grimes, Jonathan, McIntosh, Orleans, Rome Beauty, Red Delicious, Winesap and Yellow Newton — Use ½ to 1 lb. per 100 gals. water applied for full coverage. Difficult to thin Varieties such as Baldwin, Ben Davis, Duchess, Early McIntosh, Golden Delicious, Lady Apple, Northern Spy Rhode Island Greening, Steele Red, Turley, Wealthy, Yellow Transparent and York Imperial — Use 1 to 2 lbs. per 100 gals. Water. Apply in 1 spray timed between 10 and 25 days after bloom.

BEANS: Mexican Bean Beetle — 1 lb. in water to cover 1 acre. Leafhopper, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetle, Japanese Beetle — 2 lbs. in water to cover 1 acre. Lygus and Stink Bugs, Lima Bean, Pod Borer — 4 lbs. in water to cover 1 acre. Use up to 100 gals. per acre for full coverage when insects appear. Repeat at 7 to 10-day intervals if needed. Corn Earworms, Velvetbean Caterpillar — 2 to 3 lbs. in water to cover 1 acre. In low Gallonage equipment, use 50-mesh screen strainers only and provide sufficient agitation to insure suspension. lbs. per acre in water to cover. May be applied up to the day of harvest.

BLUEBERRIES: Blueberry Maggot, Cherry and Cranberry Fruitworms, European Fruit Lecanium Japanese Beetle — 3 to 4 lbs. per 100 gals. water. Apply 3 weeks before harvest and repeat 10 days later or as necessary. Apply 125 to 150 gals. diluted spray per acre.

E.P.A. Reg. No. 3743-213
E.P.A. EST. No. 3743-SC-1